

## Evaluation of pre-clinical safety and toxicology of *Althaea officinalis* extracts as naturopathic medicine for common carp (*Cyprinus carpio*)

Soleimany V.; Banaee M.\*; Mohiseni M.;  
Nematdoost Hagi B.; Mousavi Dehmourdi L.

Received: April 2015

Accepted: September 2015

### Abstract

The current study was done to investigate the preclinical safety and toxicology of *Althaea officinalis* extract as naturopathic medicine in common carp (*Cyprinus carpio*). Specimens were treated with 0 (control), 2.5, 5 and 10 g of marshmallow extract for 45 days. Plasma biochemical parameters were measured after 15 and 45 days. Total protein, albumin and globulin levels ( $p < 0.05$ ) were significantly higher in the fish fed with 10 g *A. officinalis* extract than that in control groups on day 45. Although, alanine aminotransferase (ALT), aspartate aminotransferase (AST) and alkaline phosphatase (ALP) activities significantly decreased ( $p < 0.05$ ) in fish fed with *A. officinalis* extract on day 15. The use of the *A. officinalis* extract (10 g) led to a significant increase in AST, lactate dehydrogenase (LDH) and ALP activities and cholesterol levels on day 45 ( $p < 0.05$ ) and a significant decrease in plasma glucose and cholesterol levels on day 15 ( $p < 0.05$ ). There was no significant difference in glucose levels and creatine kinase (CK) activity between all treatments and the control group on day 45 ( $p > 0.05$ ). During the experimental period, triglyceride levels noticeably decreased in fish fed with 2.5 g of *A. officinalis* extract ( $p < 0.05$ ). Although, administration of marshmallow extract up to 5 g per kg of feed did not show any side effect on fishes, the use of the *A. officinalis* extract (10 g) led to cytotoxicity and modifications in blood biochemical parameters of fish. Therefore, we recommend the use of the lower concentrations than 10 g *A. officinalis* extract in prospective clinical studies.

**Keywords:** Biochemical parameters; Common carp; Marshmallow; Medicinal plants, pre-clinical study

---

Natural Resources and the Environment Faculty, Behbahan Khatam Alanbia University of Technology, Iran

\* Corresponding author's Email: mahdibanaee@yahoo.com

## Introduction

Medical plants are used in folk veterinary medicine for treating or preventing many diseases in domesticated animals (Viegi *et al.*, 2003; Suresh Kumar and Mishra, 2004). Herbal products provide an important source of potential medicines from which humans have produced phytomedicines and herbal remedies. Herbal medicines often contain organic chemical compounds with different roles; including chemotherapeutic, immune-stimulant, bacteriostatic, bactericidal, antifungal and anti-parasitical functions (Banaee, 2010; Ahmadi *et al.*, 2012; Asadi *et al.*, 2012). In the last two decades, many studies have been conducted to determine the feasibility of using herbal medicine in prevention and curing of aquatic animals' diseases (Yin *et al.*, 2006; Divyagnaneswari *et al.*, 2007; Ardó *et al.*, 2008; Yin *et al.*, 2009).

There are several basic methods for the selection of medicinal plants including a random selection of plant species, a choice based on plants' therapeutic properties and their ethnomedical use, examining literature on phytochemical compounds and pharmacological activities as well as using chemotaxonomic approaches.

Traditionally, it is being thought that plants and plant products are non-toxic and do not have any adverse side effects, however; phytochemical ingredients of plant extracts are chemicals that are similar to those found in synthetic drugs with the same

potential to cause serious adverse effects. Therefore, choosing plant species is the first step in evaluating medicinal properties of plants and the assessment of a crude medicine is an essential part of establishing its correct identity. Prior to including any crude medicine in herbal pharmacopoeia; pharmacognostical parameters must be established. Thus, in introducing herbal medicine the pharmacognostical parameters by pre-clinical pharmacology and toxicology studies must be determined. In these studies, evaluating the changes in biochemical parameters of blood is a common clinical procedure in evaluating the medicinal effects on specific organs, especially in the case of existing difficulties in studying the changes in tissues. Measurement of biochemical markers offers comprehensive information on the physiological state of cells and tissues, particularly the liver, as a target issue. This information can be used in the pharmaceutical and toxicological evaluation to determine non-toxic doses of the drug (Banaee *et al.*, 2010).

In this study, marshmallow was chosen as a candidate drug to be used in the aquaculture industry. Marshmallow (*A. officinalis*) is a medicinal plant, the roots, leaves and flowers of which are usually used in traditional medicines in many countries all over the world (Wynn and Fougère, 2007). The flowers of marshmallow are in three colors: white, pink, and pinkish red. Sadighara *et al.* (2012) found out that

antioxidant activities and the amount of flavonoids in marshmallow extract are heavily dependent on the color of the flowers. *A. officinalis* L. are rich in mucilage polysaccharides, antioxidants, flavonoids, terpenes, and several terpenoids, sterols, saturated and unsaturated fatty acid, the amino acid asparagines, coumarins, kaempferol, phenolic acids, quercetin, sugars, tannins, and volatile oil (Elmastas *et al.*, 2004; Sartoratto *et al.*, 2004; Tešević *et al.*, 2012; Puri *et al.*, 2014). They could be found in some other medicinal plants, too. Some of these compounds such as carvacrol, thymol, linalool,  $\beta$ -bisabolene, terpinene, and 8,1- cineole, p-cymene, eugenol, furfural and camphor have antibacterial, antifungal and antiviral properties (Sartoratto *et al.*, 2004; Burt *et al.*, 2005; Jung *et al.*, 2007; Astani *et al.*, 2010; Abbaszadeh *et al.*, 2014; Bilia *et al.*, 2014; Kubiça *et al.*, 2014) or demonstrate antioxidant activity (Jun *et al.*, 2014; Puri *et al.*, 2014; Ramos *et al.*, 2014). Aristatile *et al.* (2009a;b) verified the liver-protective properties of carvacrol against the toxicity of D-glucosamine in rats. Similarly, Ezhumalai *et al.* (2014) reported that the activity of aminotransferase which had increased because of a diet rich in fats, decreased to a normal level after carvacrol administration. Considering the aforementioned properties, marshmallow may be a good alternative in the prevention and cure of bacterial, viral and fungal infections and oxidative stress.

Nevertheless, marshmallow's ingredients may have adverse side effects on the physiology and biology of fishes. Therefore, preclinical evaluation and safety evaluation of the marshmallow for fish are necessary. Hence, this study was done to collect basic information on the effects of marshmallow extract on biochemical parameters of blood in common carp.

Specimens of common carp, *C. carpio* were selected for the present study based on the following criteria: The family Cyprinidae is well represented amongst the fish inhabiting the freshwaters of Iran. This species can be found in abundant quantities throughout the year, and is of great commercial importance to people in developing countries. This species is voraciously omnivorous and is easily adapted to artificial diets. Also, this fish is hardy in nature for handling and transport. Therefore, common carp is a suitable aquatic animal for bioassay testing.

## Materials and methods

### Fish

Common carp, *C. carpio*, (with the average weight of  $37.65 \pm 4.40$  g and total length of  $14.15 \pm 0.8$  cm), were used in this study according to National Ethical Framework for Animal Research in Iran (Mobasher *et al.*, 2008). Fishes were purchased from a commercial farm in Shush, Khuzestan Province, Iran, and they were transferred to the aquaculture laboratory of Natural Resources Faculty,

Behbahan Khatam Alanbia University of Technology. Fishes were randomly distributed in 12 fiberglass tanks (200 L) and acclimatized in aerated freshwater ( $24 \pm 2^\circ\text{C}$ ; pH,  $7.4 \pm 0.2$ ; 16 L/8D; 40% water exchange rate/day) for two weeks before use. During the acclimatization period, specimens were fed two times daily with commercial diet from Beyza Feed Mill, Shiraz, Iran.

#### *Phytochemical analysis of Marshmallow (A. officinalis) flower*

*Essential oil extraction of A. officinalis*  
Marshmallow (*A. officinalis*) flower was selected from a list of herbs commonly used as medicinal plants in Iran. The hydro-distillation was done using a Clevenger-type apparatus. Dried flowers of *A. officinalis* were purchased from the local market. In the experiment, 100 g of dried flower of *A. officinalis* (as powder) was distilled with 1000 mL of water for 3 hours. The amount of the essential oil obtained was expressed in mL/100 g calculated on the basis of dry matter content.

#### *Characterization of the oil extract by Gas Chromatography (GC-MS)*

The analytical GC-MS system used was an Agilent GC-MSD system (Agilent Technologies-5975C-MS, 7890A-GC) with helium as the carrier gas at a constant linear velocity of 1 ml/min. The transfer, source and quadrupole temperatures were  $280^\circ\text{C}$ ,  $230^\circ\text{C}$  and  $150^\circ\text{C}$  respectively, operating at ionization energy of 70 eV. A HP-5MS capillary column ( $30.0\text{ m} \times 0.25\text{ mm ID}$

$\times 0.25\text{ }\mu\text{m}$  film thickness) was used and programmed from 60 to  $240^\circ\text{C}$  at 8.5/min. Oil samples (20  $\mu\text{L}$ ) were diluted with acetone (1000  $\mu\text{L}$ ). The injection volume was 0.2  $\mu\text{L}$ , the split ratio was 1:50 and the injector temperature was  $280^\circ\text{C}$ . Composition values were recorded as percentage area based on the total ion current chromatogram.

#### *Extract of A. officinalis*

The powder of dried flower *A. officinalis* was mixed with distilled water and ethanol (1:1), and the mixture was put on the shaker for 24 hours at room temperature. The resulting hydroalcoholic extract was filtered through Whatman filter paper and evaporated to dryness on a rotary evaporator until it became creamy, and was then dried in an oven at  $50^\circ\text{C}$  that finally gave 8 g (8% of initial amount) of dried powder. The concentration used in the experiment was based on the dry weight of the extract.

#### *Fish diet preparation*

The formulated fish feed was prepared in the laboratory using the powder of a commercial feed obtained from Beyza Feed Mill, Shiraz, Iran. In order to enrich the normal diet, the 2.5, 5 and 1 g of *A. officinalis* extracts were mixed with 1 kg feed powder. Each supplemented diet was mixed with distilled water (1mL/g) until a homogenous mixture was obtained. This mixture was passed through a meat grinder, producing extruded string

shapes, which were dried in an oven at 55°C for 12 h and then were broken to produce pellets approximately 10 mm in length. The pellets were packed and stored at -20°C in a freezer until used. Although no supplement was added to the feed, the control diet was prepared using the same process.

#### *The final experiment*

The final experiment was done in a completely randomized design with 3 treatments and 1 control. The fishes were fed the commercial diet enriched with the hydroalcoholic extract of flower *A. officinalis* 0.25% (2.5 g), 0.5% (5 g), and 1% (10 g) of the commercial diet and each was repeated 3 times. Common carp were fed a diet supplemented with marshmallow extract for 45 days. During experiments, specimens were monitored for appetite. Fish appetite was evaluated based on the weight of intestinal contents and their feeding behavior. After 15 and 45 days, 9 fish were captured randomly from each group and then anesthetized with clove powder solution (200 mg/L). Then, the fish blood was collected from their tail stem, and stored at 4 °C in sterilized glass vials with heparin as anticoagulant. The blood was centrifuged for 15 min at 4000 g, at 4 °C. Plasma samples were immediately stored at -21 °C until biochemical analysis.

#### *Biochemical parameters of blood*

Determination of the biochemical parameters was done using the kits supplied by Pars Azmoon Company and a UV/ VIS spectrophotometer (model UNICO 2100). Each blood biochemical parameter was determined by a certain method. Total plasma protein was measured at 540 nm by the Biuret reaction. The albumin assay is based on the dye-binding properties of plasma albumin with bromocresol green. An increase in the blue-green color was measured at 630 nm. The plasma globulin was based on the ratio of albumin versus total protein (Johnson *et al.*, 1999). Plasma glucose was measured by the glucose-oxidase method at 500 nm (Sacks, 1999). Plasma cholesterol levels by the CHOD-PAP enzymatic method at 510 nm, triglyceride levels by GPO-PAP enzymatic method at 546 nm (Rifai *et al.*, 1999) and creatinine by the JAFFE method and at 510 nm (Foster-Swanson *et al.*, 1994). The activity of aspartate aminotransferase (AST) and alanine aminotransferase (ALT) in plasma was determined by NADPH consumption and its conversion to NAD<sup>+</sup> at 340 nm. Lactate dehydrogenase (LDH) in plasma was determined based on the conversion of pyruvate to lactate at 340 nm, alkaline phosphatase (ALP) based on converting nitro phenol phosphate into nitrophenol and phosphate at 405 nm, creatinine phosphokinase (CK) based on the conversion of creatinine phosphate into creatinine at 340 nm and based on optical density (OD)

absorption and the formula presented in the kits' manual (Moss and Henderson, 1999). All biochemical parameters were determined according to the instructions provided in the kit's manual.

#### Data analysis

A significant difference in the biochemical parameters of specimens treated with the different concentrations of *A. officinalis* extracts was examined using one-way ANOVA. All the data were examined for normality (Shapiro-wilk test). Means were compared by Tukey's test and a  $p < 0.05$  was considered statistically significant. Statistical analyses were performed using SPSS (IBM, 19) software. Data are presented as mean  $\pm$  SD.

#### Results

The phytochemical analysis of the marshmallow flower sample used in this study identified 43 compounds, representing 97.823 % of the total oil content (Fig. 1 and Table 1). Table 1 represents major constituents as follows 6,10,14-trimethyl-2-Pentadecanone (47.31%), Carvacrol (17.65%), 2-Pentadecanone (11.22%), Dodecanoic acid (2.322%), n-Tetradecanoic acid (4.917%), n-Tetradecanol (1.978%), n-Nonanoic acid (1.176%), Thymol (1.073%), Methyl hexadecanoate (1.312%).

The plasma AST activities significantly decreased in fish which were fed for 15 days with feeds supplemented with 5 and 10 g of *A. officinalis* extract ( $p < 0.05$ ). The highest

AST activity was observed in fish fed with 10 g *A. officinalis* extract-supplemented feed for 45 days.

No significant difference was distinguished in AST activity after 15 days of treatment. When fish were fed with diets supplemented with 5 g *A. officinalis* extract for 45 days, the ALT activities significantly decreased ( $p < 0.05$ ).

Dietary intake of *A. officinalis* extract had significantly decreased the plasma ALP activity after 15 days of feeding ( $p < 0.05$ ). ALP activity decreased significantly in fish fed with diets supplemented with 5 g *A. officinalis* extract for 45 days ( $p < 0.05$ ).

In fish that received 2.5 g *A. officinalis* extract-supplemented feeds for 15 days, there was significant difference ( $p < 0.05$ ) in the CPK activity. However, after 45 days of feeding with extract supplemented diet, there was no significant alteration in the CPK activity.

In the present study, although no significant change in the LDH activity was observed in the fish that were fed with different doses of *A. officinalis* extract for 15 days, this activity increased significantly in fish which were fed diets supplemented with 10 g marshmallow extract for 45 days ( $p < 0.05$ ).

When administered as a diet supplement for 15 days, all the doses of the *A. officinalis* extract tested significantly decreased the plasma glucose levels ( $p < 0.05$ ). However, there was no significant difference in glucose levels between the treatments and the control group on day 45 ( $p > 0.05$ ).




Figure 1: Gas chromatography analysis of essential oil of the marshmallow (*Althaea officinalis*) flower.

No significant changes were observed in the total protein and globulin levels on day 45 and 15 ( $p>0.05$ ). However, after 15 days of feeding on *A. officinalis* extract-supplemented feed, a significant decrease was observed in plasma albumin. Fish that were fed for 45 days with 5 and 10 g of *A. officinalis* extract-supplemented diet exhibited significant increase of albumin ( $p<0.05$ ). The cholesterol significantly decreased in the fish fed for 15 days with feed

supplemented with different doses of *A. officinalis* extract ( $p<0.05$ ). However, no significant changes were observed in cholesterol levels in fish after being fed *A. officinalis* extract on day 45 ( $p>0.05$ ).

The triglyceride levels significantly decreased in fish fed with 2.5 g of *A. officinalis* extract-supplemented diet for 15 and 45 days ( $p<0.05$ ). No significant changes were observed in creatinine levels of fish ( $p>0.05$ ).

**Table 2: Enzymatic activities in plasma of common carp fed with *Althaea officinalis* extract as supplement.**

Biochemical parameters	Concentration of <i>A. officinalis</i> extract in diet (g/Kg)	Sampling time	
		15 <sup>th</sup> day	45 <sup>th</sup> day
AST (U.L <sup>-1</sup> )	Control (0.0 g)	59.89±15.45 <sup>c</sup>	38.15±10.47 <sup>ab</sup>
	2.5 g	48.97±5.04 <sup>bc</sup>	32.45±8.17 <sup>a</sup>
	5.0 g	28.61±5.23 <sup>a</sup>	54.11±12.74 <sup>bc</sup>
	10.0 g	36.06±7.09 <sup>ab</sup>	60.86±17.30 <sup>c</sup>
ALT (U.L <sup>-1</sup> )	Control (0.0 g)	11.69±2.39 <sup>a</sup>	15.55±1.86 <sup>b</sup>
	2.5 g	13.45±1.23 <sup>a</sup>	11.36±3.77 <sup>ab</sup>
	5.0 g	15.44±5.87 <sup>a</sup>	10.15±2.64 <sup>a</sup>
	10.0 g	15.11±4.53 <sup>a</sup>	15.11±2.02 <sup>b</sup>
LDH (U.L <sup>-1</sup> )	Control (0.0 g)	166.54±21.22 <sup>a</sup>	247.77±7.38 <sup>a</sup>
	2.5 g	185.24±21.55 <sup>a</sup>	210.17±46.17 <sup>a</sup>
	5.0 g	164.75±46.76 <sup>a</sup>	224.42±61.20 <sup>a</sup>
	10.0 g	157.63±19.34 <sup>a</sup>	300.12±20.94 <sup>b</sup>
ALP (U.L <sup>-1</sup> )	Control (0.0 g)	56.83±18.79 <sup>b</sup>	48.71±6.18 <sup>b</sup>
	2.5 g	27.88±2.45 <sup>a</sup>	45.80±10.95 <sup>b</sup>
	5.0 g	27.57±1.84 <sup>a</sup>	29.87±8.15 <sup>a</sup>
	10.0 g	33.70±3.11 <sup>a</sup>	58.21±8.35 <sup>b</sup>
CPK (U.L <sup>-1</sup> )	Control (0.0 g)	821.27±244.54 <sup>a</sup>	1206.93±280.03 <sup>a</sup>
	2.5 g	1432.07±269.30 <sup>b</sup>	956.11±330.41 <sup>a</sup>
	5.0 g	1191.33±578.86 <sup>ab</sup>	1200.30±293.98 <sup>a</sup>
	10.0 g	1126.44±261.23 <sup>ab</sup>	972.19±105.10 <sup>a</sup>

Significant differences between data were characterized by alphabet letters when compared with control data in the same test group along time (One way ANOVA,  $p < 0.05$ ). Values sharing the same superscript letter indicate no significant differences compared with control value during the same experiment time ( $p > 0.05$ ). Data are expressed as means ± S.D. Enzyme activity is expressed as units per liter (U.L<sup>-1</sup> or U/L).

**Table 3: Biochemical parameters in plasma of common carp fed with *Althaea officinalis* extract as supplement.**

Biochemical parameters	Concentration of <i>A. officinalis</i> extract in diet (g/Kg)	Sampling time	
		15 <sup>th</sup> day	45 <sup>th</sup> day
Glucose (mg.dL <sup>-1</sup> )	Control (0.0 g)	95.79±10.79 <sup>b</sup>	68.19±10.05 <sup>a</sup>
	2.5 g	53.65±11.35 <sup>a</sup>	56.64±10.74 <sup>a</sup>
	5.0 g	61.24±4.35 <sup>a</sup>	66.60±19.78 <sup>a</sup>
	10.0 g	59.27±9.74 <sup>a</sup>	66.86±9.34 <sup>a</sup>
Total Protein (g.dL <sup>-1</sup> )	Control (0.0 g)	3.72±0.90 <sup>a</sup>	3.70±0.63 <sup>a</sup>
	2.5 g	3.15±0.16 <sup>a</sup>	4.70±0.64 <sup>a</sup>
	5.0 g	3.41±0.55 <sup>a</sup>	4.10±0.93 <sup>a</sup>
	10.0 g	3.13±0.31 <sup>a</sup>	4.73±0.68 <sup>a</sup>

Continued Table 3:

Biochemical parameters	Concentration of <i>A. officinalis</i> extract in diet (g/Kg)	Sampling time	
		15 <sup>th</sup> day	45 <sup>th</sup> day
Albumin (g.dL <sup>-1</sup> )	Control (0.0 g)	2.24±0.31 <sup>b</sup>	1.58±0.58 <sup>a</sup>
	2.5 g	1.47±0.23 <sup>a</sup>	1.43±0.41 <sup>a</sup>
	5.0 g	1.67±0.41 <sup>a</sup>	2.73±0.38 <sup>b</sup>
	10.0 g	1.69±0.31 <sup>a</sup>	3.60±0.49 <sup>c</sup>
Globulin (g.dL <sup>-1</sup> )	Control (0.0 g)	1.48±0.68 <sup>a</sup>	2.12±1.12 <sup>ab</sup>
	2.5 g	1.68±0.26 <sup>a</sup>	3.28±0.30 <sup>b</sup>
	5.0 g	1.74±0.83 <sup>a</sup>	1.37±1.19 <sup>a</sup>
	10.0 g	1.45±0.46 <sup>a</sup>	1.13±0.76 <sup>a</sup>
Cholesterol (mg.dL <sup>-1</sup> )	Control (0.0 g)	59.03±2.37 <sup>c</sup>	73.04±14.35 <sup>ab</sup>
	2.5 g	49.86±4.45 <sup>b</sup>	66.37±9.76 <sup>a</sup>
	5.0 g	36.69±6.69 <sup>a</sup>	67.20±16.57 <sup>a</sup>
	10.0 g	35.52±7.12 <sup>a</sup>	99.05±24.76 <sup>b</sup>
Triglyceride (mg.dL <sup>-1</sup> )	Control (0.0 g)	239.44±59.28 <sup>b</sup>	261.74±25.08 <sup>b</sup>
	2.5 g	153.05±13.05 <sup>a</sup>	180.05±65.39 <sup>a</sup>
	5.0 g	192.49±43.84 <sup>ab</sup>	223.47±35.49 <sup>ab</sup>
	10.0 g	208.45±23.63 <sup>ab</sup>	261.50±63.47 <sup>b</sup>
Creatinine (mg.dL <sup>-1</sup> )	Control (0.0 g)	0.57±0.26 <sup>a</sup>	0.30±0.06 <sup>a</sup>
	2.5 g	0.45±0.22 <sup>a</sup>	0.35±0.05 <sup>a</sup>
	5.0 g	0.38±0.13 <sup>a</sup>	0.30±0.06 <sup>a</sup>
	10.0 g	0.36±0.19 <sup>a</sup>	0.31±0.09 <sup>a</sup>

Significant differences between data were characterized by alphabet letters when compared with control data in the same test group along time (One way ANOVA,  $p < 0.05$ ). Values sharing the same superscript letter indicate no significant differences compared with control value during the same experiment time ( $p > 0.05$ ). Data are expressed as means±S.D.

## Discussion

Dodecanoic acid and n-Tetradecanoic acid are the major saturated fatty acid found in the essential oil of *A. officinalis*. Monoterpenes have various biological properties including anticancer (Astani *et al.*, 2010), antimicrobial, antifungal, anti-genotoxic, anti-inflammatory, insecticidal and antioxidant (Aydmn and Türkez, 2014). *P-Cymene* (1-isopropyl-4-methylbenzene) is a natural hydrocarbon that is a component of

essential oils, which are natural products extracted from marshmallow. *P-Cymene* has anti-bacterial properties. Borneol and camphor are bicyclic monoterpenes found in marshmallow extract.  $\alpha$ -Terpineol (TPN), a volatile monoterpene alcohol, is relatively non-toxic and one of the major components of the essential oils of marshmallow. Benzaldehyde, n-decanal, is mainly used as a food and flavoring additive and can be found in many fruit and plants. Menthol is a cyclic terpene

alcohol detected in marshmallow extract. Therefore, marshmallow has the potential to be used for prevention and treatment of bacterial, viral, and fungal infections.

During the experimental periods, no mortalities or changes in the appetite of the fish were observed. Platel *et al.* (2002) evidenced the favorable effect of medicinal plants on digestion and a stimulating effect on bile secretion and the activity of pancreatic enzymes. Moreover, adding plants extracts to the diet can affect the ability of the fish to find food by stimulating their sense of smell and can encourage them to eat more (Adams, 2005).

In this study, oral administration of the *A. officinalis* extract significantly decreased the plasma ALT activity at a dose of 5 g compared to the control group on day 45 ( $p < 0.05$ ). Supplementation of *A. officinalis* extract to fish significantly decreased the activity of alkaline phosphatase on day 15 ( $p < 0.05$ ). Treatment of the fish with 2.5 g *A. officinalis* extract decreased activities AST, and LDH in plasma compared to the control group. Although, increased creatine kinase (CK) activity was observed in the plasma of fish fed 2.5 g *A. officinalis* extract on day 15 ( $p < 0.05$ ), CK activity returned to the normal levels in fish treated with *A. officinalis* extract at the end of the experimental period (Table 2). More hydrophilic flavonoids interacting at the membrane surface through hydrogen bonding; may reduce the access of oxidants and per-oxidants,

thus protecting the structure and function of cellular membranes (Oteiza *et al.*, 2005). Oral administration of thymol and carvacrol activated antioxidant enzymes and protected liver cells against severe damages (Hashemipour *et al.*, 2013). Carvacrol supplement reduced AST, ALT, and LDH activity in plasma of the fish treated with D-glucosamine (Jayakumar *et al.*, 2012). The effects of marshmallow extract on the activity of plasma enzymes can be compared with the effects of other extracts such as *Allium cepa* and *A. sativum* (Al-Salahy, 2002), *Curcuma longa* (Deshpande *et al.*, 2003) *Cyperus rotundus* (Suresh Kumar and Mishra, 2004), *Pterocarpus santalinus* (Palanisamy *et al.*, 2007), *Aloe vera*, *Clematis hirsute*, *Cucumis prophetarum* (Alqasoumi *et al.*, 2008), *Hybanthus enneaspermus* (Premalakshmi and Thenmozhi, 2011). The useful effect of *A. officinalis* in decreasing ALP activity might be due to the presence of minerals, particularly calcium levels in the *A. officinalis* extract. A low concentration of silymarin (Banaee *et al.*, 2011), yarrow extract (Nafisi Bahabadi *et al.*, 2014) and garlic (Al-Salahy, 2002) in diet of fish are proven to regulate the plasma activities of AST, ALT, ALP, CK and LDH.

The results show that AST and LDH activities increased in plasma of fish fed a diet supplemented with 10 g/Kg of *A. officinalis* extract on day 45 which may be due to leakage of these enzymes from the liver cytosol into blood. Some

monoterpenes such as d-limonene,  $\alpha$ -pinene, myrcene and linalool, stylosin could have cytotoxic effects (Aydın and Türkez, 2014). *P-Cymene* is metabolized by the liver and the major *p*-cymene metabolite is cuminyl alcohol which can cause mitochondrial toxicity by affecting the process of energy production in hepatocytes (Custódio *et al.*, 2011). Toxicological studies show that safranal, and furfural, are other toxic compounds found in plants such as marshmallow. Dimethyl glutarate is an ester that may cause olfactory toxicity (Morris *et al.*, 1991). Furfural turns into the toxic pyromucic acid in liver after oxidation. Therefore, furfural may indirectly cause oxidative stress and damage the liver cells (Veićković *et al.*, 2011). Increased activity of AST, ALT and ALP was reported in the rats treated with furfural during 7 days (Veićković *et al.*, 2011).

A significant lower concentration of glucose was noted in fish fed a diet enriched with *A. officinalis* extract as compared to control fishes on day 15 ( $p < 0.05$ ). The probable mechanism of *A. officinalis* extract may be partly due to the stimulation of insulin secretion or may be attributed to the activation of glycogen synthesis and healthy hepatic function (Ji *et al.*, 2007). Moreover, this was probably because of the flavonoids in the plants which reduce glucose uptake in the intestine via the inhibition of sodium-dependent glucose transport (Song *et al.*, 2002). Decreased glucose levels in the blood of rainbow trout fed with silymarin extract (Banaee *et al.*,

2011), yarrow extract (Nafisi Bahabadi *et al.*, 2014) and African catfish fed with onion and garlic extracts are reported (Al-Salahy, 2002).

Since there is a close relationship between the rate of protein synthesis in liver and total protein concentration in plasma (Banaee *et al.*, 2011), the increased concentration of total protein in plasma of the fish fed with marshmallow extract may indicate increased protein synthesis rate in the liver. Amino acids present in the *A. officinalis* extract may increase protein synthesis in the liver and other tissues. Although, there was no significant difference in plasma globulin level on days 15, oral administration of 2.5 g *A. officinalis* extract significantly increased plasma globulin level on day 45 ( $p < 0.05$ ). Oral administration of marshmallow extract decreased albumin level on day 15, while the albumin level in the fish fed with marshmallow extract (5 and 10 g) increased significantly on day 45 compared with the control group ( $p < 0.05$ ). An increased level of total protein, albumin and globulin in plasma of the fish fed with diets enriched with *Echinacea purpurea* and *Silybum marianum* was reported by Bohlouli Oskoi *et al.* (2012) and Banaee *et al.* (2011). No significant changes were reported in the levels of albumin and total protein in plasma of fish fed with diets enriched with onion and garlic extract (Al-Salahy, 2002).

Supplementation of *A. officinalis* extract in the diet significantly reduced

the cholesterol levels in the plasma ( $p < 0.05$ ). Interference with cholesterol absorption from intestine due to administration of *A. officinalis* might have played a role in decreasing plasma cholesterol levels. The beneficial effects of *A. officinalis* on cholesterol might be due to pectin and mucilage (Ezhumalai *et al.*, 2014) and phytosterols (Marinangeli *et al.*, 2006; Jones *et al.*, 2007). Triglycerides or triacylglycerol are neutral fats, major energy reserves for the body stored in the adipose tissue. Therefore, decrease in triglyceride levels in plasma of fish fed with 2.5 g *A. officinalis* extract in diet may be due to increased lipogenesis and decrease lipolysis and ketogenesis. In the lipogenesis process, triglycerides and fatty acids are synthesized and stored in adipose tissue. Decreases in cholesterol and triglyceride levels were also reported in blood of rainbow trout and catfish respectively fed with silymarin extract (Banaee *et al.*, 2011), yarrow extracts (Nafisi Bahabadi *et al.*, 2014) and onion and garlic extract (Al-Salahy, 2002).

This work has demonstrated that *A. officinalis* is by far relatively non-toxic (2.5 and 5 g) in terms of biochemical parameters. Other findings obtained in the present study are that *A. officinalis* extract exhibited moderate cytotoxicity in high dose (10 g), while its extract has moderate antioxidant properties at doses of 2.5 and 5 g. Indeed, the potential in the plant to influence several clinically important parameters could be linked to various levels of

pharmacological advantages. So, we recommend the use of these dosages in prospective clinical study to further evaluate the efficacy and safety of the *A. officinalis* extract as naturopathic medicine for fishes.

### Acknowledgments

The authors gratefully acknowledge the support offered (8/1/T/244) from the Behbahan Khatam Alanbia University of Technology. Also, the authors are grateful to Maryam Banaee, our English editor, for proofreading the manuscript.

### References

- Abbaszadeh, S., Sharifzadeh, A., Shokri, H., Khosravi, A.R. and Abbaszadeh, A., 2014. Antifungal efficacy of thymol, carvacol, eugenol and menthol as alternative agents to control the growth of food-relevant fungi. *Journal of Medical Mycology*, 24(2), 51-56.
- Adams, C., 2005. Nutrition-based health. *Feed internat*, 2, 25-28.
- Ahmadi, K., Banaee, M., Vosoghei, A.R., Mirvaghefi, A.R. and Ataimehr, B., 2012. Evaluation of the immunomodulatory effects of silymarin extract (*Silybum marianum*) on some immune parameters of rainbow trout, *Oncorhynchus mykiss* (Actinopterygii, Salmoniformes). *Acta Ichthyologica Et Piscatoria*, 42 (2), 113-120.
- Alqasoumi, S.I., Al-Howiriny, T.A. and Abdel-Kader, M.S., 2008. Evaluation of the hepatoprotective

- effect of *Aloe vera*, *Clematis hirsute*, *Cucumis prophetarum* and bee propolis against experimentally induced liver injury in rats. *International Journal of Pharmacology*, 4(3), 213-217.
- Al-Salahy, M.B., 2002.** Some physiological studies on the effect of onion and garlic juices on the fish, *Clarias lazera*. *Fish Physiology and Biochemistry*, 27, 129-142.
- Ardó, L., Yin, G., Xu, P., Váradi, L., Szigeti, G., Jeney, Z. and Jeney, G., 2008.** Chinese herbs (*Astragalus membranaceus* and *Lonicera japonica*) and boron enhance the non-specific immune response of Nile tilapia (*Oreochromis niloticus*) and resistance against *Aeromonas hydrophila*. *Aquaculture*, 275, 26-33.
- Aristatile, B., Al-Numair, K.S., Veeramani, C. and Pugalendi, K.V., 2009a.** Effect of CVL on hepatic marker enzymes and antioxidant status in D-galactosamine-induced hepatotoxicity in rats. *Fundamental Clinical Pharmacology*, 23, 757-765.
- Aristatile, B., Al-Numair, K.S., Veeramani, C. and Pugalendi, K.V., 2009b.** Antihyperlipidemic effect of carvacrol on D-galactosamine-induced hepatotoxic rats. *Journal of Basic and Clinical Physiology and Pharmacology*, 20(1), 15-27.
- Asadi, M.S., Mirvaghefi, A.R., Nematollahi, M.A., Banaee, M. and Ahmadi, K., 2012.** Effects of Watercress (*Nasturtium nasturtium*) extract on some immunological parameters of rainbow trout (*Oncorhynchus mykiss*). *Open Veterinary Journal*, 2(1), 32-39.
- Astani, A., Reichling, J. and Schnisler, P., 2010.** Comparative study on the antiviral activity of selected monoterpenes derived from essential oils. *Phytotherapy Research*, 24(5), 673-679.
- Aydın, E. and Türkez, H., 2014.** In vitro cytotoxicity, genotoxicity and antioxidant potentials of thymol on human blood cells. *Journal of Essential Oil Research*, 26(2), 133-140.
- Banaee, M., 2010.** Influence of silymarin in decline of sublethal diazinon-induced oxidative stress in rainbow trout (*Oncorhynchus mykiss*). Ph.D. Thesis, Aquaculture and Environmental Department, Natural Resource Faculty, Natural Resource and Agriculture Collage, Tehran University, Iran. 149 P.
- Banaee, M., Sureda, A., Mirvaghefi, A.R. and Rafei, G.R., 2011.** Effects of long-term silymarin oral supplementation on the blood biochemical profile of rainbow trout (*Oncorhynchus mykiss*). *Fish Physiology and Biochemistry*, 37, 887-896.
- Bilia, A.B., Santomauro, F., Sacco, C., Bergonzi, M.C. and Donata, R., 2014.** Essential oil of *Artemisia annua* L., An extraordinary component with numerous antimicrobial properties. Hindawi

- Publishing Corporation. *Evidence-Based Complementary and Alternative Medicine*. ID: 159819. 7 P.
- Bohlouli Oskoi, S., Tahmasebi Kohyani, A., Salati, A.P., Parseh, A. and Sadeghi, E., 2012.** Effects of dietary administration of *Echinacea purpurea* on growth indices and biochemical and hematological indices in rainbow trout (*Oncorhynchus mykiss*) fingerlings. *Fish Physiology and Biochemistry*, 38, 1029-1034.
- Burt, S.A., Vlieland, R., Haagsman, H.P., Edwin, E.J. and Veldhuizen, A., 2005.** Increase in activity of essential oil components carvacrol and thymol against *Escherichia coli* O157:H7 by addition of food stabilizers. *Journal of Food Protection*, 68(5), 919-926.
- Custódio, J.B.A., Ribeiro, M.V., Silva, F.S.G., Machado, M. and Sousa, M.C., 2011.** The essential oils component *p*-cymene induces proton leak through Fo-ATP synthase and uncoupling of mitochondrial respiration. *Journal of Experimental Pharmacology*, 3, 69-76.
- Deshpande, U.R., Joseph, L.J. and Samuel, A.M., 2003.** Hepatobiliary clearance of labelled Mebrofenin in normal and D-GalN Hcl induced hepatitis rats and the protective effect of turmeric extract. *Indian Journal of Physiology and Pharmacology*, 47(3), 332-336.
- Divyagnaneswari, M., Christyapita, D. and Dinakaran, M.R., 2007.** Enhancement of nonspecific immunity and disease resistance in *Oreochromis mossambicus* by *Solanum trilobatum* leaf fractions. *Fish & Shellfish Immunology*, 23, 249-259.
- Elmastas, M., Ozturk, L., Gokce, I., Erenler, R. and Aboul-Enein, H.Y., 2004.** Determination of antioxidant activity of marshmallow flower (*Althaea officinalis* L.). *Bioanalytical*, 37(9), 1859-1869.
- Ezhumalai, M., Radhiga, T. and Pugalendi, K.V., 2014.** Antihyperglycemic effect of carvacrol in combination with rosiglitazone in high-fat diet-induced type 2 diabetic C57BL/6J mice. *Molecular and Cellular Biochemistry*, 385(1-2), 23-31.
- Foster-Swanson, A., Swartzentruber, M. and Roberts, P., 1994.** Reference interval studies of the rate-blanked creatinine, Jaffe method on BM /Hitachi Systems in Six U.S. Laboratories (Abstract). *Clin. Chem.* No. 361.
- Hashemipour, H., Kermanshahi, H., Golian, A. and Veldkamp, T., 2013.** Effect of thymol and carvacrol feed supplementation on performance, antioxidant enzyme activities, fatty acid composition, digestive enzyme activities, and immune response in broiler chickens. *Poultry Science*, 92(8), 2059-2069.

- Jayakumar, S., Madankumar, A., Asokkumar, S., Raqhunandhakumar, S., Gokuladhas, K., Kamaraj, S., Divya, M.G. and Devaki, T., 2012.** Potential preventive effect of carvacrol against diethylnitrosamine-induced hepatocellular carcinoma in rats. *Molecular and Cellular Biochemistry*, 360(1), 51-60.
- Ji, S.C., Jronh, G.S., Gwang-Soon, I.M., Lee, S.W., Yoo, J.H. and Takii, K., 2007.** Dietary medicinal herbs improve growth performance, fatty acid utilization, and stress recovery of Japanese flounder. *Fisheries Science*, 73(1), 70-76.
- Johnson, A.M., Rohlf, E.M. and Silverman, L.M., 1999.** Proteins. In: Burtis CA, Ashwood ER, editors. *Tietz Textbook of Clinical Chemistry. 3rd ed. W.B. Saunders Company, Philadelphia.* pp. 477-540.
- Jones, P.J.H., Demonty, I., Chan, Y., Herzog, Y. and Pelled, D., 2007.** Fish-oil esters of plant sterols differ from vegetable-oil sterol esters in triglycerides lowering, carotenoid bioavailability and impact on plasminogen activator inhibitor-1 (PAI-1) concentrations in hypercholesterolemic subjects. *Lipids Health Disease*, 6, 28.
- Jun, H.J., Lee, J.H., Kim, J., Jia, Y., Kim, K.H., Hwang, K.Y., Yun, E.J., Do, K.R. and Lee, S.L., 2014.** Linalool is a PPAR $\alpha$  ligand that reduces plasma TG levels and rewires the hepatic transcriptome and plasma metabolome. *Journal of Lipid Research*, 55, 1098-1110.
- Jung, K.H., Yoo, S.K., Moon, S.K. and Lee, U.S., 2007.** Furfural from pine needle extract inhibits the growth of a plant pathogenic fungus, *Aternaria mali*. *Mycobiology*, 35(1), 39-43.
- Kubiça, T.F., Alves, S.H., Weiblen, R. and Lovato, L., 2014.** In vitro inhibition of the bovine viral diarrhea virus by the essential oil of *Ocimum basilicum* (basil) and monoterpenes. *Brazilian Journal of Microbiology*, 45(1), 209-214.
- Marinangeli, C.P., Varady, K.A. and Jones, P.J., 2006.** Plant sterols combined with exercise for the treatment of hypercholesterolemia, overview of independent and synergistic mechanisms of action. *The Journal of Nutritional Biochemistry*, 17, 217-224.
- Mobasher, M., Aramesh, K., Aldavoud, S.J., Ashrafganjooei, N., Divsalar, K. and Phillips, C.J.C., 2008.** Proposing a national ethical framework for animal research in Iran. *Iranian Journal Public Health*, 37(1), 39-46.
- Morris, J.B., Clay, R.J., Trela, B.A. and Bogdanffy, M.S., 1991.** Deposition of dibasic esters in the upper respiratory tract of the male and female Sprague-Dawley rat. *Toxicology and Applied Pharmacology*, 108(3), 538-546.
- Moss, D.V. and Henderson, A.R., 1999.** Clinical enzymology. In: Burtis CA, Ashwood ER, editors.

- Tietz Textbook of Clinical Chemistry. 3rd ed. W.B. Saunders Company, Philadelphia. pp. 617-721.
- Nafisi Bahabadi, M., Banaee, M., Taghian, M. and Nematdoust Hagi, B., 2014.** Effects of dietary administration of yarrow extract on growth performance and blood biochemical parameters of rainbow trout (*Oncorhynchus mykiss*). *International Journal of Aquatic Biology*, 2(5), 275-285.
- Oteiza, P.I., Erlejman, A.G., Verstraeten, S.V., Keen, C.L. and Fraga, C.G., 2005.** Flavonoid-membrane interactions, A protective role of flavonoids at the membrane surface? *Clinical & Developmental Immunology*, 12(1), 19-25.
- Palanisamy, D., Syamala, S. and Bhojraj, S., 2007.** Protective and therapeutic effects of the Indian medicinal plant *Pterocarpus santalinus* on D-galactosamine-induced liver damage. *Asian Journal of Traditional Medicines*, 2(2), 51-57.
- Platel, K., Rao, A., Saraswahi, G. and Srinivasan, K., 2002.** Digestive stimulant action of three Indian spice mixes in experimental rats. *Die Nahrung*, 46, 394-398.
- Premalakashmi, V. and Thenmozhi, C.D., 2011.** Hepatoprotective effect of hydroethanolic extract of *Hybanthus enneaspermus* against paracetamol induced liver damage in rats. *International Journal of Biotechnology Biochemistry*, 7(2), 239-244.
- Puri, A, Srivastava, P., Pandey, P., Yadav, R.S. and Bhatt, P.C., 2014.** Scopolamine induced behavioral and biochemical modifications and protective effect of *Celastrus paniculatus* and *Angelica glauca* in rats. *International Journal of Nutrition, Pharmacology, Neurological Diseases*, 4(3), 158-169.
- Ramos, M., Beltrán, A., Peltzer, M., Valente, A.J. and del Carmen Garrigós, M., 2014.** Release and antioxidant activity of carvacrol and thymol from polypropylene active packaging films. *LWT-Food Science and Technology*, 58(2), 470-477.
- Rifai, N., Bachorik, P.S. and Albers, J.J., 1999.** Lipids, lipoproteins and apolipoproteins. In: Burtis CA, Ashwood ER, eds. *Tietz Textbook of Clinical Chemistry. 3rd ed. W.B. Saunders Company, Philadelphia.* pp. 809-861.
- Sacks, D.B., 1999.** Carbohydrates. In: Burtis CA, Ashwood ER, eds. *Tietz Textbook of clinical chemistry. 3rd ed. W.B. Saunders Company, Philadelphia.* pp. 766-85.
- Sadighara, P., Gharibi, S., Moghadam Jafari, A., Jahed Khaniki, G.R. and Salari, S., 2012.** The antioxidant and flavonoids contents of *Althaea officinalis* L. flowers based on their color. *Avicenna Journal of Phytomedicine*, 2(3), 113-117.

- Sartoratto, A., Machado, A.L., Delarmelina, C., Figueira, G.M., Duarte, M.C. and Rehder, V.L., 2004.** Composition and antimicrobial activity of essential oils from aromatic plants used in Brazil. *Brazilian Journal of Microbiology*, 35, 275-280.
- Song, J., Kwon, O., Chen, S., Daruwala, R., Eck, P. and Park, J.B., 2002.** Flavonoid inhibition of sodium-dependent vitamin C transport 1 (SVCT 1) and glucose transporter isoform 2 (GLUT 2), intestinal transporters for vitamin C and glucose. *JBC*, 277, 15252-60.
- Suresh Kumar, S.V. and Mishra, S.H., 2004.** Protective effect of rhizome extract of *Cyperus rotundus* Linn against thioacetamide induced hepatotoxicity in rats. *Indian Drugs*, 41(3), 165-169.
- Tešević, V., Vajs, V., Lekić, S., Dordević, I., Novaković, M., Vujisić, L. and Todosijević, M., 2012.** Lipid composition and antioxidant activities of the seed oil from three Malvaceae species. *Archives of Biological Science Belgrade*, 64(1), 221-227.
- Veičković, D., Milenković, S. and Stojanović, D., 2011.** Enzymochemical and biochemical changes in the liver of rats induced by furfural. *Acta Medica Mediane*, 50(2), 34-38.
- Viegi, L., Pieroni, A., Guarrera, P.M. and Vangelisti, R., 2003.** A review of plants used in folk veterinary medicine in Italy as basis for a databank. *Journal of Ethnopharmacology*, 89, 221-244.
- Wynn, S.G., and B.J. Fougère., 2007.** Veterinary herbal medicine. *Mosby Elsevier, St. Louis, Missouri, USA*. 736 P.
- Yin, G., Ardó, L., Thompson, K.D., Adams, A. and Jeney, G., 2009.** Chinese herbs (*Astragalus radix* and *Ganoderma lucidum*) enhance immune response of carp, *C. carpio*, and protection against *Aeromonas hydrophila*. *Fish & Shellfish Immunology*, 26(1), 140-145.
- Yin, G., Jeney, G., Racz, T., Xu, P., Jun, X. and Jeney, Z., 2006.** Effect of two Chinese herbs (*Astragalus radix* and *Scutellaria radix*) on non-specific immune response of tilapia, *Oreochromis niloticus*. *Aquaculture*, 253 (1-4), 39-47.