

## **Evaluation the performance of anaesthetic effects of Tobacco and Ketamine on Grass Carp, *Ctenopharyngodon idella* (Valenciennes,1844)**

**Khara H.<sup>1\*</sup>; Ahmadnezhad M.<sup>2</sup>; Rahbar M.<sup>3</sup>; Khodadust A.<sup>3</sup>**

Received: October 2018

Accepted: February 2019

### **Abstract**

The present study aimed to determine the optimal concentration of three anaesthetics, name MS222, ketamine, and tobacco extract, in grass carp juveniles ( $107.78 \pm 22.11$  g body weight and  $226.74 \pm 13.06$  mm total body length). Treatments were as follows:  $100 \text{ mg L}^{-1}$  of MS222 (control group), 5.5, 8, 11, 18.5, and  $22 \text{ mg L}^{-1}$  of ketamine, and 10, 13.33, 20, 26.27, and  $40 \text{ mg L}^{-1}$  of tobacco extract. In addition, changes in blood and biochemical parameters were measured at different concentrations. Considering the anaesthesia induction and recovery durations times, the optimal concentration for anaesthesia of studied substances was obtained as follows:  $100 \text{ mg L}^{-1}$  of MS222 ( $139.00 \pm 20.00$  seconds),  $18.5 \text{ mg L}^{-1}$  of ketamine ( $140.10 \pm 20.20$  seconds), and  $40 \text{ mg L}^{-1}$  of tobacco extract ( $174.00 \pm 20.09$  seconds). Based on the results, the shortest recovery time from anaesthesia was observed at  $26.27 \text{ mg L}^{-1}$  of tobacco extract ( $870.00 \pm 5.3$  seconds) and  $8 \text{ mg L}^{-1}$  of ketamine ( $967.00 \pm 5.5$  seconds), respectively. The results indicated that the anaesthesia induction and recovery duration was dependent on the concentration of anaesthetics. Moreover, significant differences were observed between the three anaesthetics in terms of blood parameters. The study findings also revealed a significant difference between treatments in biochemical parameters (cortisol, glucose, and lactate).

**Keywords:** Anaesthetics, Blood, Grass Carp, *Ctenopharyngodon idella*, MS<sub>222</sub>, Ketamine, Tobacco

---

1-Department of Fisheries, Lahijan Branch, Islamic Azad University, Lahijan, Iran

2-Inland Waters Aquaculture Research Center, Iranian Fisheries Sciences Research Institute, Agricultural Research, Education and Extension Organization (AREEO), Bandar Anzali, Iran

3-Young Researchers and Elite Club, Lahijan Branch, Islamic Azad University, Lahijan, Iran

\*Corresponding author's Email: h.khara1974@yahoo.com

## Introduction

In the modern aquaculture, handling practices like transportation, sizing, grading, weighing, stripping and blood collection are stressful for fish. To reduce the effects of stress, some chemical substances and anaesthetics are routinely utilized for fish before handling (Ross and Ross, 1999; Wagner *et al.*, 2003).

Anaesthetic agents are potentially used for light sedation and non-invasive procedures to fulfil anaesthesia in order to avoid inflicting pain during surgery and interventions (Ross and Ross, 2008; Neiffer and Stamper, 2009). Many anaesthetics agents have been used in aquaculture practices, including tricaine methanesulfonate (MS-222), clove oil, xylocaine, benzocaine, 2-phenoxyethanol, quinaldine, quinaldine sulphate, and metomidate (Gutierrez and Herrera, 1995; Masee *et al.*, 1995; Ortuno *et al.*, 2002; Small, 2003; Kiessling *et al.*, 2009).

However, some parameters such as efficacy, cost effectiveness, availability and safety have been considered for a valuable agent. Although, ketamine and tobacco extracts have been experimentally have been applied in researchs (Di Marco *et al.*, 2011; Mohammadi and Khara, 2015; Dinesh *et al.*, 2017) but, the use of these agents and their impacts on physiological parameters remain unexplored, since no maximum residue levels and species-specific dosage have been determined. At the present, only MS-222 is approved as the most widely used anaesthetics chemical in fish (Topic

Popovic *et al.*, 2012; Chambel *et al.*, 2015).

The evaluation of fish health following induced anaesthesia is linked to study of blood biochemical properties. It has been reported that, anaesthetics may potentially alter blood biochemistry in many fish species (Iwama *et al.*, 1989). The stress-induced alterations of post-anaesthesia in blood chemistry indices occur within seconds or minutes after fish are treated (Mazeaud and Mazeaud, 1981; Gingerich and Drottar, 1989). Therefore, precautions must be taken into an account to ensure that handling procedures do not change the indices of interest. These changes are attributed to occur in plasma hormones, energy metabolism and electrolytes balance that enable biologists to use aforementioned parameters for evaluation of fish stress responses and health status (Wagner and Congleton, 2004).

Some studies have been found that MS-222 or clove oil affect blood biochemical parameters, including glucose (Davis and Griffin, 2004), cortisol (Wagner *et al.*, 2003), total protein (Cataldi *et al.*, 1998) and amino acid (Morales *et al.*, 1990). These previous studies mainly focused on conventional anaesthetic agents, while little attention was considered to the physiological effects plant based or other anaesthetic agents. Hence, understanding of the new anaesthetic agents effects have a great practical application in aquaculture. In the current study, we tested haematological profiles and blood biochemical

parameters to determine the potential stress responses of two anaesthetics; ketamine and tobacco extract to compare with MS-222 as conventionally approved anaesthetic on juvenile grass carp. Therefore, the objectives of this study were to determine the appropriate anaesthetic concentrations of ketamine and tobacco compared to MS-222 as a potential substitute anaesthetics and the effect of anaesthetics on blood and biochemical parameters to identify the best anaesthetic for Grass Carp juveniles, particularly regard to reducing stress.

## **Materials and methods**

### *Fish*

165 pieces of Grass Carp juveniles ( $107.78 \pm 22.11$  g body weight and  $226.74 \pm 13.06$  mm total body length) were provided from local hatchery near to experiment place. The study was conducted at the Lahijan Branch of Islamic Azad University (Iran). Before beginning of the experiment, fish were initially acclimatized for 10 days in one circular fiberglass tank (1500 L) in a flow-through water system. Water temperature of the holding tanks was maintained at 18-21 °C throughout the experiment. Fish were not fed for 24 h before anaesthesia treatments and blood sampling (Ross and Ross, 2008).

### *Anaesthesia and experimental procedure*

MS-222 was purchased from Sigma-Aldrich Chemicals Ltd (USA, Saint Louis). Ketamine was supplied by Rotexdemica Company (Dresden, Germany) and tobacco extract was

provided from Tobacco Company in Rasht (Iran). Before using, the clove oil was dissolved in 96% ethanol (1: 9 v/v) and MS-222 in water to prepare a stock solution (Chambel *et al.*, 2015). Preliminary trials of ethanol used in each trial did not have any noticeable effects on the fish. The applied concentrations of tobacco extract were calculated as 50% 24 h LC<sub>50</sub> (24 h LC<sub>50</sub> of tobacco extract on Grass Carp obtained from preliminary test). Thus, 100 mg of tobacco extract were measured and mixed with 1 L of water to prepare stock solution of 100 mg L<sup>-1</sup> concentration (Agokei and Adebisi, 2010; Dinesh *et al.*, 2017). Ketamine was dissolved in 1 L of water to give desired concentrations (Di Marco *et al.*, 2011).

In this study, 11 groups of 15 Grass carp (MS-222: 1 Treatment, 15 pieces, tobacco: 5 Treatments, 75 pieces, ketamine: 5 Treatments, 75 pieces) each were compared in this study:

Control: based on the previous studies, the exact concentrations of MS-222 was served as control treatment to compare effectiveness of other anaesthetics as alternative for anaesthetizing Grass carp (Topic Popovic *et al.*, 2012; Chambel *et al.*, 2015).

Two groups with blood sampling immediately after anaesthesia in each concentration of tobacco and following recovery time courses.

Two groups blood sampling immediately after anaesthesia in each concentration of ketamine and following recovery time courses.

Three replicates for each treatment were designed for each group, each

held in 30 L glass aquaria (38cm length×27cm width×47cm height) containing freshwater plus the anaesthetic agents. Induction time of anaesthesia was recorded from adding the anaesthetic agent to the water till no opercula movements, decreased respiratory rate and weak response to strong tactile stimuli (Phase II) have been observed.

After anaesthesia, to evaluate the haematological and blood biochemistry parameters, samples were taken from caudal severance and divided into two portions. Half of each blood sample was immediately transferred to heparinized tubes for haematological examination and the rest was put in non-heparinized tubes for serum analysis. For biochemical assays, blood samples were immediately centrifuged (3000 g for 10 min) at room temperature and then the serum was separated and stored at -20°C until analysis. Haematological parameters, including erythrocyte count (RBC), haemoglobin value (Hb, g dl<sup>-1</sup>), haematocrit value (Hct, %), leukocyte count (WBC), mean corpuscular volume (MCV, fl), mean corpuscular haemoglobin (MCH, pg), and mean corpuscular haemoglobin concentration (MCHC, %) have been measured by using standard procedure (Harikrishnan *et al.*, 2003).

Differential leukocyte counts (neutrophil, lymphocyte, monocyte and eosinophil) were determined by Giemsa staining method of blood smears using a light microscope. The smears obtained from heparinized sample were first air-dried, fixed in 96 % ethanol for

30 min and stained in Giemsa to determine differential leukocyte counts (Klontz, 1994).

Radioimmunoassay (RIA) was used for measurement of cortisol. An assay kit of cortisol was purchased from Immunotech Company, France. Commercial assay kit was contained two necessary reagents, i.e., antibody and tracer (labelled antigen). Aliquots containing antibody and tracer were transferred to assay tubes (polystyrene tube). Tracer and antibody were added in a further 100 µl of assay buffer, and the tubes were left to incubate overnight at 4 °C. The content of polystyrene tubes (antibody, tracer, and serum sample) was discharged completely and then polystyrene tubes applied to a Gamma counter (Wallac, Finland) for measurement of absorption spectrum. Subsequently hormone concentration was calculated using related standard curve. The concentration of glucose and lactate were measured spectrophotometrically using kits supplied by Pars Azmun Diagnostics, Tehran, Iran.

#### *Statistical analysis*

A Kruskal–Wallis test was used to determine the difference between the induction and recovery times of different concentrations of the each anaesthetic agent (Differences among means of blood indices were analyzed by one-way ANOVA followed by Duncan's new multiple range tests). Differences were considered significant at  $p < 0.05$ . Data were analyzed using SPSS statistical package (SPSS,

Chicago IL, USA) and expressed as mean±standard error of the mean (SE).

## Results

Significant differences ( $p<0.05$ ) in the induction times of anaesthesia and recovery times were observed at different concentrations of the anaesthetic agents. Induction time of anaesthesia was shortest and longest at MS-222 and 5.5 mg L<sup>-1</sup> of ketamine, respectively (Table 1). Furthermore, the shortest and longest recovery times were observed in tobacco extract and ketamine at concentrations of 26 mg L<sup>-1</sup> and 18.5 mg L<sup>-1</sup>, respectively (Table 2). Haematological parameters of Grass carp including WBC, RBC, Ht and Hb were significantly influenced by anaesthetic agents in induction and recovery times (Tables 2 and 3). There were dose-dependent variations in WBC, RBC, Ht and Hb levels in induction times among different concentrations anaesthetic agents (Table 2). In induction times of anaesthesia and recovery times, significant differences were observed in WBC, RBC, Hb, Ht and MCV values

among different concentrations of anaesthetic agents. The highest levels of RBC, WBC and Ht in induction times were observed in tobacco at concentrations of 10, 13 and 40 mg L<sup>-1</sup> (Table 2). Whereas, in recovery times, the highest levels of RBC and Hb values were obtained in ketamine treatment at concentrations of 5.5 mg L<sup>-1</sup> (Table 3).

Moreover, in induction and recovery times of Grass Carp significant changes in differential leukocyte counts (lymphocyte, neutrophil and monocyte) among different concentrations of anaesthetic agents were detected (Tables 4 and 5).

In the induction and recovery times, serum cortisol concentration was considerably higher in tobacco extract and ketamine at concentrations of 10 mg L<sup>-1</sup> and 22 mg L<sup>-1</sup>, respectively (Fig. 1). A similar result was also seen in the serum glucose concentration (Fig. 2). Lactate concentration in the induction and recovery times was significantly higher in ketamine at concentrations of 5.5 mg L<sup>-1</sup> and 22 mg L<sup>-1</sup>, respectively (Fig. 3).

**Table 1: Induction and recovery times (s) of juvenile grass carp anesthetized with MS-222 (control), Ketamine and Tobacco as anaesthetic agents.**

Anesthetics concentration	Induction anesthesia time (second)	Recovery time (second)
MS 222 (100 mg L <sup>-1</sup> )	139 ± 20.1	4100 ± 5.3
Ketamine (5.5 mg L <sup>-1</sup> )	289 ± 10	2112 ± 20
Ketamine (8 mg L <sup>-1</sup> )	243± 15.1	967 ± 5.5
Ketamine (11 mg L <sup>-1</sup> )	243± 20	2595 ± 30
Ketamine (18.5 mg L <sup>-1</sup> )	140 ± 10.1	4776 ± 20.2
Ketamine (22 mg L <sup>-1</sup> )	144 ± 6	4353 ± 16
Tobacco (10 mg L <sup>-1</sup> )	175 ± 10.2	1500 ± 7.5
Tobacco (13 mg L <sup>-1</sup> )	175± 10.1	1521± 15.3
Tobacco (20 mg L <sup>-1</sup> )	174± 10.4	1498± 30
Tobacco (26 mg L <sup>-1</sup> )	175± 10.5	870± 5.3
Tobacco (40 mg L <sup>-1</sup> )	174± 15	1600± 20.9

**Table 2: Effects of MS-222 (control), Ketamine and Tobacco extract as anaesthetic agents on haematocrit values of juvenile grass carp following anaesthesia.**

Anaesthetics	Concentrations	WBC ( $\times 10^3 \text{ mm}^{-3}$ )	RBC ( $\times 10^6 \text{ mm}^{-3}$ )	Hb (gr dl <sup>-1</sup> )	Ht (%)	MCV ( $\mu\text{m}^{-3}$ )	MCH (pg)	MCHC (%)
MS 222	(100 mg L <sup>-1</sup> )	4100 ± 100 <sup>f</sup>	184 ± 45.9 <sup>b</sup>	8.5 ± 0.60 <sup>a</sup>	35.6 ± 2.50 <sup>a</sup>	187 ± 6.2 <sup>b</sup>	46 ± 1	24 ± 1.5
Ketamine	(5.5 mg L <sup>-1</sup> )	4500 ± 500 <sup>de</sup>	185 ± 44.9 <sup>b</sup>	8.5 ± 0.50 <sup>a</sup>	35 ± 1 <sup>a</sup>	189 ± 1 <sup>ab</sup>	45 ± 2	23 ± 1.1
Ketamine	(8 mg L <sup>-1</sup> )	4816 ± 200 <sup>cd</sup>	155 ± 85.1 <sup>g</sup>	7.1 ± 0.10 <sup>bc</sup>	29 ± 1 <sup>cd</sup>	186 ± 0.57 <sup>b</sup>	46 ± 1	24 ± 1
Ketamine	(11 mg L <sup>-1</sup> )	4850 ± 200 <sup>cd</sup>	164 ± 31.8 <sup>f</sup>	7 ± 0.0 <sup>bc</sup>	27 ± 0.0 <sup>de</sup>	181 ± 1.7 <sup>cd</sup>	46 ± 1	25 ± 2
Ketamine	(18.5 mg L <sup>-1</sup> )	4700 ± 200 <sup>cd</sup>	173 ± 75.3 <sup>d</sup>	6.9 ± 0.10 <sup>cd</sup>	26 ± 1.5 <sup>ef</sup>	179 ± 00 <sup>d</sup>	45 ± 2.5	23 ± 0.57
Ketamine	(22 mg L <sup>-1</sup> )	5200 ± 200 <sup>c</sup>	173 ± 65.2 <sup>d</sup>	6.8 ± 0.10 <sup>cd</sup>	26 ± 0.57 <sup>ef</sup>	177 ± 00 <sup>df</sup>	45 ± 2.5	25 ± 2
Tobacco	(10 mg L <sup>-1</sup> )	6000 ± 200 <sup>b</sup>	150 ± 43.5 <sup>h</sup>	6.4 ± 0.25 <sup>d</sup>	26.4 ± 0.11 <sup>d</sup>	173 ± 2.8 <sup>f</sup>	42 ± 2.3	24 ± 0.57
Tobacco	(13 mg L <sup>-1</sup> )	4093 ± 310 <sup>g</sup>	155 ± 35 <sup>g</sup>	6.9 ± 0.15 <sup>bc</sup>	26.9 ± 0.25 <sup>bc</sup>	186 ± 2.5 <sup>b</sup>	46 ± 1	24 ± 0.57
Tobacco	(20 mg L <sup>-1</sup> )	8306 ± 390 <sup>a</sup>	149 ± 81.7 <sup>i</sup>	6.9 ± 0.20 <sup>cd</sup>	28.3 ± 0.57 <sup>cde</sup>	188 ± 1.5 <sup>bc</sup>	46 ± 0.57	25 ± 0.00
Tobacco	(26 mg L <sup>-1</sup> )	5816 ± 275 <sup>b</sup>	189 ± 35 <sup>a</sup>	8.4 ± 0.05 <sup>a</sup>	35.3 ± 0.67 <sup>a</sup>	185 ± 1 <sup>a</sup>	44 ± 0.57	24 ± 0.57
Tobacco	(40 mg L <sup>-1</sup> )	4713 ± 280 <sup>cd</sup>	160 ± 39.7 <sup>f</sup>	7.9 ± 0.05 <sup>b</sup>	30.6 ± 2.5 <sup>bc</sup>	193 ± 2.6 <sup>a</sup>	45 ± 0.57	23 ± 00

**Table 3: Effects of MS-222 (control), Ketamine and Tobacco as anaesthetic agents on haematocrit values of juvenile grass carp following recovery.**


Anaesthetics	Concentrations	WBC ( $\times 10^3 \text{ mm}^{-3}$ )	RBC ( $\times 10^6 \text{ mm}^{-3}$ )	Hb (gr dl <sup>-1</sup> )	Ht (%)	MCV ( $\mu\text{m}^{-3}$ )	MCH (pg)	MCHC (%)
MS 222	(100 mg L <sup>-1</sup> )	4100 ± 100 <sup>f</sup>	184 ± 45.9 <sup>b</sup>	8.5 ± 0.60 <sup>a</sup>	35.6 ± 2.50 <sup>a</sup>	187 ± 6.2 <sup>b</sup>	46 ± 1	24 ± 1.5
Ketamine	(5.5 mg L <sup>-1</sup> )	4500 ± 500 <sup>de</sup>	185 ± 44.9 <sup>b</sup>	8.5 ± 0.50 <sup>a</sup>	35 ± 1 <sup>a</sup>	189 ± 1 <sup>ab</sup>	45 ± 2	23 ± 1.1
Ketamine	(8 mg L <sup>-1</sup> )	4816 ± 200 <sup>cd</sup>	155 ± 85.1 <sup>g</sup>	7.1 ± 0.10 <sup>bc</sup>	29 ± 1 <sup>cd</sup>	186 ± 0.57 <sup>b</sup>	46 ± 1	24 ± 1
Ketamine	(11 mg L <sup>-1</sup> )	4850 ± 200 <sup>cd</sup>	164 ± 31.8 <sup>f</sup>	7 ± 0.0 <sup>bc</sup>	27 ± 0.0 <sup>de</sup>	181 ± 1.7 <sup>cd</sup>	46 ± 1	25 ± 2
Ketamine	(18.5 mg L <sup>-1</sup> )	4700 ± 200 <sup>cd</sup>	173 ± 75.3 <sup>d</sup>	6.9 ± 0.10 <sup>cd</sup>	26 ± 1.5 <sup>ef</sup>	179 ± 00 <sup>d</sup>	45 ± 2.5	23 ± 0.57
Ketamine	(22 mg L <sup>-1</sup> )	5200 ± 200 <sup>c</sup>	173 ± 65.2 <sup>d</sup>	6.8 ± 0.10 <sup>cd</sup>	26 ± 0.57 <sup>ef</sup>	177 ± 00 <sup>df</sup>	45 ± 2.5	25 ± 2
Tobacco	(10 mg L <sup>-1</sup> )	6000 ± 200 <sup>b</sup>	150 ± 43.5 <sup>h</sup>	6.4 ± 0.25 <sup>d</sup>	26.4 ± 0.11 <sup>d</sup>	173 ± 2.8 <sup>f</sup>	42 ± 2.3	24 ± 0.57
Tobacco	(13 mg L <sup>-1</sup> )	4093 ± 310 <sup>g</sup>	155 ± 35 <sup>g</sup>	6.9 ± 0.15 <sup>bc</sup>	26.9 ± 0.25 <sup>bc</sup>	186 ± 2.5 <sup>b</sup>	46 ± 1	24 ± 0.57
Tobacco	(20 mg L <sup>-1</sup> )	8306 ± 390 <sup>a</sup>	149 ± 81.7 <sup>i</sup>	6.9 ± 0.20 <sup>cd</sup>	28.3 ± 0.57 <sup>cde</sup>	188 ± 1.5 <sup>bc</sup>	46 ± 0.57	25 ± 0.00
Tobacco	(26 mg L <sup>-1</sup> )	5816 ± 275 <sup>b</sup>	189 ± 35 <sup>a</sup>	8.4 ± 0.05 <sup>a</sup>	35.3 ± 0.67 <sup>a</sup>	185 ± 1 <sup>a</sup>	44 ± 0.57	24 ± 0.57
Tobacco	(40 mg L <sup>-1</sup> )	4713 ± 280 <sup>cd</sup>	160 ± 39.7 <sup>f</sup>	7.9 ± 0.05 <sup>b</sup>	30.6 ± 2.5 <sup>bc</sup>	193 ± 2.6 <sup>a</sup>	45 ± 0.57	23 ± 00

**Table 4: Effects of MS-222 (control), Ketamine and Tobacco as anaesthetic agents on differential leukocyte counts of juvenile grass carp following anaesthesia.**


Anaesthetics	Concentrations	Lymphocyte (%)	Neutrophil (%)	Eosinophil (%)	Monocyte (%)
MS 222	(100 mg L <sup>-1</sup> )	74 ± 2 <sup>bcdef</sup>	20.6 ± 1.1 <sup>cd</sup>	0.33 ± 0.57	5 ± 1 <sup>ab</sup>
Ketamine	(5.5 mg L <sup>-1</sup> )	74 ± 1 <sup>bcdef</sup>	21.3 ± 0.57 <sup>cd</sup>	0.33 ± 0.57	4.3 ± 0.57 <sup>bc</sup>
Ketamine	(8 mg L <sup>-1</sup> )	75 ± 1 <sup>bcdef</sup>	22 ± 1 <sup>bc</sup>	0.33 ± 0.57	2.3 ± 0.57 <sup>abc</sup>
Ketamine	(11 mg L <sup>-1</sup> )	72 ± 1.5 <sup>bcde</sup>	23 ± 0.57 <sup>ab</sup>	0.33 ± 0.57	3.6 ± 0.57 <sup>de</sup>
Ketamine	(18.5 mg L <sup>-1</sup> )	71 ± 0.57 <sup>efg</sup>	24 ± 1 <sup>a</sup>	0.33 ± 0.57	4 ± 1 <sup>abcd</sup>
Ketamine	(22 mg L <sup>-1</sup> )	76 ± 0.57 <sup>fg</sup>	20 ± 1 <sup>cd</sup>	0.33 ± 0.57	3 ± 0 <sup>abcd</sup>
Tobacco	(10 mg L <sup>-1</sup> )	71 ± 1 <sup>b</sup>	24 ± 0.57 <sup>a</sup>	0 ± 0 <sup>d</sup>	4.6 ± 0.57 <sup>abc</sup>
Tobacco	(13 mg L <sup>-1</sup> )	73 ± 2.3 <sup>f</sup>	21.3 ± 1.5 <sup>cd</sup>	0 ± 0 <sup>bc</sup>	5 ± 1 <sup>ab</sup>
Tobacco	(20 mg L <sup>-1</sup> )	76 ± 1.5 <sup>cdefg</sup>	19.6 ± 1.5 <sup>d</sup>	0.66 ± 0.57	3.3 ± 0.57 <sup>bcd</sup>
Tobacco	(26 mg L <sup>-1</sup> )	75 ± 0.57 <sup>bcd</sup>	19.3 ± 0.57 <sup>d</sup>	0.66 ± 0.57	4.6 ± 1.4 <sup>abc</sup>
Tobacco	(40 mg L <sup>-1</sup> )	80 ± 2 <sup>a</sup>	17.3 ± 2 <sup>e</sup>	0.33 ± 0.57	1.6 ± 0.57 <sup>e</sup>

**Table 5: Effects of MS-222 (control), Ketamine and Tobacco as anaesthetic agents on differential leukocyte counts of juvenile grass carp following recovery.**


Anaesthetics	Concentrations	Lymphocyte (%)	Neutrophil (%)	Eosinophil (%)	Monocyte (%)
MS 222	(100 mg L <sup>-1</sup> )	73 ± 1.5 <sup>b</sup>	21 ± 1 <sup>ef</sup>	0.33 ± 0.57	5 ± 1.7 <sup>bc</sup>
Ketamine	(5.5 mg L <sup>-1</sup> )	76 ± 0.57 <sup>a</sup>	18.3 ± 0.57 <sup>cd</sup>	0.66 ± 0.57	4.3 ± 0.57 <sup>b</sup>
Ketamine	(8 mg L <sup>-1</sup> )	70 ± 0.57 <sup>cd</sup>	23.6 ± 0.57 <sup>a</sup>	1.3 ± 0.57	4.6 ± 0.57 <sup>bc</sup>
Ketamine	(11 mg L <sup>-1</sup> )	67 ± 1 <sup>e</sup>	28 ± 0 <sup>a</sup>	1 ± 1	4.6 ± 0.57 <sup>bc</sup>
Ketamine	(18.5 mg L <sup>-1</sup> )	67 ± 0.57 <sup>e</sup>	27 ± 0.57 <sup>bc</sup>	0.66 ± 1.1	5 ± 0 <sup>bc</sup>
Ketamine	(22 mg L <sup>-1</sup> )	68 ± 1 <sup>de</sup>	24 ± 1 <sup>ab</sup>	1.33 ± 0.57	6 ± 1 <sup>a</sup>
Tobacco	(10 mg L <sup>-1</sup> )	66 ± 2 <sup>e</sup>	26 ± 0.57 <sup>de</sup>	0.66 ± 0.57	6 ± 1 <sup>a</sup>
Tobacco	(13 mg L <sup>-1</sup> )	72 ± 2.5 <sup>bc</sup>	22 ± 0.57 <sup>a</sup>	1.33 ± 0.27	4.3 ± 0.57 <sup>b</sup>
Tobacco	(20 mg L <sup>-1</sup> )	66 ± 1.5 <sup>e</sup>	28 ± 2 <sup>a</sup>	0.66 ± 0.57	4.6 ± 0.57 <sup>bc</sup>
Tobacco	(26 mg L <sup>-1</sup> )	72 ± 2 <sup>bc</sup>	22 ± 1 <sup>cde</sup>	0 ± 0	4.6 ± 0.57 <sup>bc</sup>
Tobacco	(40 mg L <sup>-1</sup> )	72 ± 2.5 <sup>bc</sup>	22 ± 2.5 <sup>de</sup>	0.66 ± 0.57	4.3 ± 0.57 <sup>b</sup>



**Figure 1: Changes in serum cortisol concentration of juvenile grass carp following anaesthesia with MS-222 (control), ketamine and tobacco and recovery time. Data are expressed as Mean±SD; n=10 fish per group. Different letters indicate differences among groups at  $p < 0.05$ .**



**Figure 2: Changes in serum glucose concentration of juvenile grass carp following anaesthesia with MS-222 (control), ketamine and tobacco and recovery time. Data are expressed as Mean±SD; n=10 fish per group. Different letters indicate differences among groups at  $p < 0.05$ .**



**Figure 3: Changes in serum lactate concentration of juvenile grass carp following anaesthesia with MS-222 (control), ketamine and tobacco and recovery time. Data are expressed as Mean±SD; n=10 fish per group. Different letters indicate differences among groups at  $p < 0.05$ .**



## Discussion

Applications of anaesthetic agents are common practice to handle fish stress and prevent physical injuries during all aquaculture practices (Iversen *et al.*, 2003). Despite utilization of various anaesthetics agents in modern aquaculture, the knowledge of comparative effects of different suitable dosage for handling operations is still limited (Di Marco *et al.*, 1999; Fleming *et al.*, 2003; Gomulka *et al.*, 2008). Therefore, it is important to determine the appropriate anaesthetic concentrations and exposure the time to minimize the stress (Feng *et al.*, 2011). It has been recommended, the adequate induction time of an anaesthetic for fish should be below 15 min, with a recovery time of lower than 5 min (Marking and Meyer, 1985). However, Cardenas *et al.* (2016), have had been pointed that the anaesthetic efficacy of anaesthetic agents is obviously species-specific, size-dependent, and affected by water temperature.

The anaesthetic concentrations used in this study were safe and effective in reducing anesthetizing induction time in fish. The lowest anaesthetic induction time in each of the MS<sub>222</sub> test materials with a time of 139.00±20.00 seconds, followed by a concentration of 18.5 mg L<sup>-1</sup> of ketamine with a time of 140.120 20.20 sec, and 40 mg L<sup>-1</sup> tobacco with a time of 17.09±20.09 seconds.

Anaesthetic dosages used in this study were safe and effective in order to rapid induction of anaesthesia in grass carp with MS-222 followed by ketamine with the lowest concentration. A similar results on sturgeon hybrid

*Acipenser naccarii*×*Acipenser baerii* was reported in concentration of 150 mg L<sup>-1</sup> MS-222 (Di Marco *et al.*, 2011). The results of the present study demonstrated that the most effective anaesthetic that induce anaesthesia with the lowest recovery time were MS- 222, 5.5 mg L<sup>-1</sup> and 26 mg mg L<sup>-1</sup> of tobacco for Grass carp. Similarly, MS-222 efficacy for induction (≤3min) and recovery (≤5min) times were reported at the same dose of the present study (100 mg L<sup>-1</sup>) in several fish (Hseu *et al.*, 1998; Bystriansky *et al.*, 2006; Ibarra-Zatarain *et al.*, 2011). Conversely, the appropriate ranges of ketamine and tobacco extract concentrations in fish are rarely studied. However, it has been reported that the effects of ketamine and tobacco extract in terms of acquiring optimal concentration in induction and recovery times for rainbow trout was varied (Mohammadi and Khara, 2015). This discrepancy is likely due to the differences in experimental protocols or fish species studied. In line with this, Williams *et al.* (1988) have noted that there were interspecies differences in the successful dosage of anaesthetic. In this study, the minimum return time of anesthesia in Amor fish was 26.67 mg L<sup>-1</sup> of tobacco with 870.00±3.5 g followed by a concentration of 8 mg L<sup>-1</sup> of ketamine with 967±5.5 seconds. In the present study, anaesthetic induction time significantly decreased with anesthetic concentration increase.

The reaction of fish exposed to excess of the recommended dose of any anaesthetic will be accompanied with an obvious stress response. For

instance, Molinero and Gonzalez (1995) reported that gilthead sea bream *Sparus aurata* exposed to MS-222 and 2-phenoxyethanol at a dose exceeding 25 mg L<sup>-1</sup> and 0.075 mg L<sup>-1</sup> respectively, represented a stress response.

In this regard, blood chemistry parameters may help to provide some indicators to determine the optimal concentration range for anaesthesia and decrease stress (Czesny *et al.*, 2003; Feng *et al.*, 2011). Generally, cortisol and glucose levels are the main biomarkers of physiological stress that have been used as indicators (Wagner *et al.*, 2002; Gesto *et al.*, 2015). Blood erythrocytes were also used as an indicator of anaesthetic stress (Gontijo *et al.*, 2003). However, almost no study has reported that other blood variables could be used as indicators of anaesthetic stress for fish. In the present study, significant changes in plasma parameters including cortisol, glucose, and lactate levels in induction and recovery times were dose-dependent. However, the literature suggests that both glucose and lactate levels are typically correlated with an increase in cortisol (Wendelaar Bonga, 1997; Barton, 2002), as it was occurred in the present study. An increase in plasma glucose after only a few minutes of anaesthesia was observed in juvenile salmonids exposed to a lethal dose of MS-222. However, earlier studies reported that plasma glucose was stable for 12-15 minutes in salmonids exposed to 80-100 mg L<sup>-1</sup> neutralized or un-neutralized MS-222 (Houston *et al.*, 1971; Nieminen *et al.*, 1982;

Congleton, 2006). On the other hand, elevation of plasma glucose have been reported for several species of marine fish exposed to 80-100 mg L<sup>-1</sup> MS-222 (Bourne, 1984; Thomas and Robertson, 1991). It is not known whether the increase of plasma glucose observed in the present study was related to the examined species, or to the high concentration of anaesthetics used. Our results for lactate level are in accordance with the reports of other authors, who reported that lactate concentration increased significantly in all anaesthetized fish as a metabolic responses to anaesthesia (Di Marco *et al.*, 2011; Cardenas *et al.*, 2016). It has been shown that, plasma lactate level could be dependent upon secondary effects of anaesthesia, related with oxygen insufficiency for cell aerobic metabolism (Iversen *et al.*, 2003). Although, longer periods of anaesthesia or even delayed sampling points are the main necessary factors for existing of variations in this plasma metabolite.

Differential leukocyte counts are important characteristics of the health status of the fish and in many cases they are also used to evaluate the effect of drugs and anaesthetics on fish (Gomulka *et al.*, 2008; Witeska *et al.*, 2015). Very few data on the effects of anaesthetics on thrombocyte counts in fish are available. In the present study, juveniles of Grass carp subjected to different anaesthetics agents have been showed significant changes in lymphocyte and neutrophil counts. In line with our findings, Witeska *et al.* (2015) had reported an increase in neutrophil and monocyte of Grass carp

under handling anaesthesia with 2-phenoxyethanol or etomidate.

Several studies have been shown that control of the stress responses in fish is dependent on the type of anaesthetic, the concentration used and exposure time (Strange and Schreck, 1978; Barton and Peter, 1982; Davidson *et al.*, 2000). In the present research, the appropriate anaesthetic concentrations for juvenile of Grass Carp were just relative values, which should be concluded based on blood biochemical parameters, anaesthetic time, recovery time, exposure time, and so on. According to the present study, juvenile of Grass Carp were physiologically affected by MS-222 as same as tobacco, indicating that tobacco extract is an appropriate alternative to MS-222. Since, tobacco extract is inexpensive, easily obtained, and non-toxic to fish and human (Dinesh *et al.*, 2017), it can be potentially used as a proper anaesthetic for fish.

In general, data analysis showed that the physiological effects were higher with tobacco anaesthetics than ketamine and MS222. The results of current and previous studies had indicated the importance of considering all the physiological potential or other anaesthetic effects of other substances used in aquaculture and fishery studies. It is clear that various anaesthetics can have a definite effect on the fish's blood-type indices. Therefore, when choosing an anaesthetic agents for fish whose anaesthetic effects on them are unknown, a careful examination should be made, especially when changes in

the blood-counting index are used as the decisive indicator in a study.

### **Acknowledgments**

The present work was conducted at the Lahijan Islamic Azad University, Iran. The authors wish to thank the manager and staff of laboratory center for their valuable contribution.

### **References**

- Agokei, O.E. and Adebisi, A.A., 2010.** Tobacco as anaesthetic for fish handling procedures. *Journal of Medicinal Plants Research*, 4(14), 1396-1399.
- Barton, B.A. and Peter, R.E., 1982.** Plasma cortisol stress response in fingerling rainbow trout, *Salmo gairdneri* Richardson, to various transport conditions, anaesthesia, and cold shock. *Journal of Fish Biology*, 20, 39–51.
- Barton, B.A., 2002.** Stress in fishes: a diversity of responses with particular reference to changes in circulating corticosteroids. *Integrative and Comparative Biology*, 42, 517–525.
- Bourne, P.K., 1984.** The use of MS 222 (tricaine methanesulphonate) as an anaesthetic for routine blood sampling in three species of marine teleosts. *Aquaculture*, 36, 313-321.
- Bystriansky, J.S., LeBlanc, P.J. and Ballantyne, J.S., 2006.** Anaesthetisation of Arctic charr *Salvelinus alpinus* (L) with tricaine methanesulphonate or 2-phenoxyethanil for immediate blood sampling. *Journal of Fish Biology*, 69, 613-621.

- Cardenas, C., Toni, C., Martos-Sitcha, J.A., Cardenas, S., de las Heras, V., Baldisserotto, B., Heinzmann, B.M., Vazquez, R. and Mancera, J.M., 2016.** Effects of clove oil, essential oil of *Lippia alba* and 2-phe anaesthesia on juvenile meagre, *Argyrosomus regius* (Asso, 1801). *Journal of Applied Ichthyology*, 32, 693–700.
- Cataldi, E., Di Marco, P., Mandich, A. and Cataudella, S., 1998.** Serum parameters of Adriatic sturgeon *Acipenser naccarii* (Pisces: Acipenseriformes): Effects of temperature and stress. *Comparative Biochemistry and Physiology A*, 121, 351–354.
- Chambel, J., Pinho, R., Sousa, R., Ferreira, T., Baptista, T., Severiano, V., Mendes, S. and Pedrosa, R., 2015.** The efficacy of MS-222 as anaesthetic agent in four freshwater aquarium fish species. *Aquaculture Research*, 46(7), 1582–1589.
- Congleton, J.L., 2006.** Stability of some commonly measured blood-chemistry variables in juvenile salmonids exposed to a lethal dose of the anaesthetic MS-222. *Aquaculture Research*, 37, 1146–1149.
- Czesny, S., Rinchar, J., Garcia, M.A. and Dabrowski, K.D., 2003.** The effect of fasting, prolonged swimming, and predator presence on energy utilization and stress in juvenile walleye (*Stizostedion vitreum*). *Physiology and Behavior*, 79, 597–603.
- Davidson, G.W., Davie, P.S., Young, G. and Fowler, R.T., 2000.** Physiological responses of rainbow trout *Oncorhynchus mykiss* to crowding and anesthesia with AQUIS. *Journal of the World Aquaculture Society*, 31, 105–114.
- Davis, K.B. and Griffin, B.R., 2004.** Physiological responses of hybrid striped bass under sedation by several anesthetics. *Aquaculture*, 233, 531–548.
- Di Marco, P., McKenzie, D.J., Mandich, A., Bronzi, P., Castaldi, E. and Cataudella, S., 1999.** Influence of sampling conditions on blood chemistry values for Adriatic sturgeon *Acipenser naccarii* (Bonaparte, 1836). *Journal of Applied Ichthyology*, 15, 73–77.
- Di Marco, P., Petochi, T., Longobardi, A., Priori, A., Finoi, M.G., Donadelli, V., Corsalini, I. and Marino, G., 2011.** Efficacy of tricaine methanesulphonate, clove oil and medetomidine-ketamine and their side effects on the physiology of sturgeon hybrid *Acipenser naccarii* × *Acipenser baerii*. *Journal of Applied Ichthyology*, 27, 611–617.
- Dinesh, R., Prakash, C., Poojary, N., Kantharajan, G. and Abraham, S., 2017.** Tobacco (*Nicotiana tabacum*) - A novel and futuristic sedative for fish transport in India. *International Journal of Fisheries and Aquatic Studies*, 5(3), 369–371.
- Feng, G., Zhuang, P., Zhang, L., Kynard, B., Shi, X., Duan, M., Liu, J. and Huang, X., 2011.** Effect of anaesthetics MS-222 and clove oil on blood biochemical parameters of juvenile Siberian sturgeon

- (*Acipenser baerii*). *Journal of Applied Ichthyology*, 27, 595–599
- Fleming, G.J., Heard, D.J., Floyd, R.F. and Riggs, A., 2003.** Evaluation of propofol and medetomidine-ketamine for short-term immobilization of Gulf of Mexico sturgeon (*Acipenser oxyrinchus desotii*). *Journal of Zoo and Wildlife Medicine*, 34, 153–158.
- Gesto, M., Hernández, J., López-Patiño, M.A., Soengas, J.L. and Míguez, J.M., 2015.** Is gill cortisol concentration a good acute stress indicator in fish? A study in rainbow trout and zebrafish. *Comparative Biochemistry and Physiology, Part A*, 188, 65–69.
- Gingerich, W.H. and Drottler, K.R., 1989.** Plasma catecholamine concentrations in rainbow trout (*Salmo gairdneri*) at rest and after anesthesia and surgery. *General and Comparative Endocrinology*, 73, 390–397.
- Gomulka, P., Wlasow, T., Velisek, J., Svobodová, Z. and Chmielinska, E., 2008.** Effects of eugenol and MS-222 anaesthesia on Siberian sturgeon *Acipenser baerii* Brandt. *Acta Veterinaria Brno*, 77, 447–453.
- Gontijo, A.M., Barreto, R.E., Speit, G., Reyes, V.A.V., Volpato, G.L. and Salvadori, D.M.F., 2003.** Anesthesia of fish with benzocaine does not interfere with comet assay results. *Mutation Research*, 534, 165–172.
- Gutierrez, M.R. and Herrera, A.E., 1995.** Evaluation of the repeated use of xylocaine as anesthetic for the handling of breeding carp (*Cyprinus carpio*). *Aquaculture*, 129, 431–436.
- Harikrishnan, R., Harikrishnan, M. and Balasundaram, C., 2003.** hematological and biochemical parameters in common Carp, *Cyprinus carpio*, following herbal treatment for *Aeromonas hydrophila* infection. *Aquaculture*, 221(1-4), 41–50.
- Houston, A.H., Madden, J.A., Woods, R.J. and Miles, H.M., 1971.** Some physiological effects of handling and tricaine methanesulphonate anesthetization upon the brook trout, *Salvelinus fontinalis*. *Journal of Fisheries Research Board of Canada*, 28, 625–633.
- Hseu, J.R., Yeh, S.L., Chu, Y.T. and Ting, Y.Y., 1998.** Comparison of efficacy of five anesthetic goldlined sea bream, *Sparus sarba*. *Acta Zoologica Taiwanica*, 9, 35–41.
- Ibarra-Zatarain Z., Ibarra-Castro, L., Alvarez-Lajonchère, L., Aguilar, N.G. and Sánchez-Téllez, J.L., 2011.** The use of three anaesthetics for handling spotted rose snapper *Lutjanus guttatus* (Pisces, Lutjanidae) broodstock. *Revista de Biología Marina y Oceanografía*, 46, 471–476.
- Iversen, M., Finstad, B., MacKinley, R.S. and Eliassen, R.A., 2003.** The efficacy of metomidate, clove oil, Aqui-STM and Benzoak as anaesthetics in Atlantic salmon (*Salmo salar* L.) smolts, and their potential stress-reducing capacity. *Aquaculture*, 221, 549–566.
- Iwama, G.K., McGeer, J.C. and Pawluk, M.P., 1989.** The effects of

- five fish anaesthetics on acid-base balance, haematocrit, blood gases, cortisol, and adrenaline in rainbow trout. *Canadian Journal of Zoology*, 67, 2065–2073.
- Kiessling, A., Johansson, D., Zahl, I.H. and Samuelsen, O.B., 2009.** Pharmacokinetics, plasma cortisol and effectiveness of benzocaine, MS-222 and isoeugenol measured in individual dorsal aorta-cannulated Atlantic salmon (*Salmo salar*) following bath administration. *Aquaculture*, 286, 301–308.
- Klontz, G.W., 1994.** Fish hematology. In: Techniques in Fish Immunology, Stolen, J.S., T.C. Flecher, A.F. Rowley, T.C. Zelikoff, S.L. Kaattari and S.A. Smith (Eds.). Vol. 2, SOS Publications, USA, ISBN: 0962550582. pp. 121-132.
- Marking, L.L. and Meyer, F.P., 1985.** Are better anesthetics needed in fisheries? *Fisheries*, 10, 2–5.
- Massee, K.C., Rust, M.B., Hardy, R.W. and Stichney, R.R., 1995.** The effectiveness of tricaine, quinaldine sulfate and metomidate as anesthetics for larval fish. *Aquaculture*, 134, 351–359.
- Mazeaud, M.M. and Mazeaud, F., 1981.** Adrenergic responses to stress in fish. In: Stress and Fish (ed. by A.D. Pickering), Academic Press, New York, USA. pp. 49-75.
- Mohammadi, M. and Khara, H., 2015.** Effect of different anesthetic agents (clove oil, tricaine methanesulfonate, ketamine, tobacco) on hematological parameters and stress indicators of rainbow trout *Oncorhynchus mykiss*, Walbaum, 1792. *Comparative Clinical Pathology*, 24(5), 1039–1044.
- Molinero, A. and Gonzalez, J., 1995.** Comparative effects of MS 222 and 2-phenoxyethanol on gilthead sea bream (*Sparus aurata* L.) during confinement. *Comparative Biochemistry and Physiology*, 111A, 405–414.
- Morales, A.E., Garcia-Rejon, L. and Dela Higuera, M., 1990.** Influence of handling and/or anaesthesia on stress response in rainbow trout. Effects on liver primary metabolism. *Comparative Biochemistry and Physiology*, 95A, 87-93.
- Neiffer, D.L. and Stamper, M.A., 2009.** Fish sedation, anesthesia, analgesia, and euthanasia: Considerations, methods, and types of drugs. *Institute for Laboratory Animal Research*, 50, 343–360.
- Nieminen, M., Laitnen, M. and Pasanen, P., 1982.** Effects of anaesthesia with tricaine (MS 222) on the blood composition of the splake (*Salvelinus fontinalis* X *Salvelinus namaycush*). *Comparative Biochemistry and Physiology*, 73C, 271-276.
- Ortuno, J., Esteban, M.A. and Meseguer, J., 2002.** Effects of four anaesthetics on the innate immune response of gilthead seabream (*Sparus aurata* L.). *Fish and Shellfish Immunology*, 12, 49–59.
- Ross, L.G. and Ross, B., 1999.** Anaesthetic and sedative techniques for aquatic animals second edition. Blackwell Science Ltd., Oxford. 159 P.

- Ross, L.G. and Ross, B., 2008.** Anaesthetic and sedative techniques for aquatic animals. Blackwell, Oxford. 240 P.
- Small, B.C., 2003.** Anesthetic efficacy of metomidate and comparison of plasma cortisol responses to tricaine methanesulfonate, quinaldine and clove oil anesthetized channel catfish *Ictalurus punctatus*. *Aquaculture*, 218, 177–185.
- Strange, R.J. and Schreck, C.B., 1978.** Anesthetic and handling stress on survival and cortisol concentration in yearling chinook salmon (*Oncorhynchus tshawytscha*). *Journal of the Fisheries Research Board of Canada*, 35, 345–349.
- Svobodov, A.Z., Groch, L., Flajshans, M., Vykusov, A.B. and Machov, A.J., 1997.** The effect of long-term therapeutic bath of malachite green on common carp (*Cyprinus carpio* L.). *Acta Veterinaria Brno*, 66, 111–117.
- Thomas, P. and Robertson, L., 1991.** Plasma cortisol and glucose stress responses of red drum (*Sciaenops ocellatus*) to handling and shallow water stressors and anesthesia with MS-222, quinaldine sulfate and metomidate. *Aquaculture*, 96, 69–86.
- Topic Popovic, N., Strunjak-Perovic, I., Coz-Rakovac, R., Barisic, J., Jadan, M., Persin Berakovic, A. and Sauerborn Klobucar, R., 2012.** Tricaine methane-sulfonate (MS-222) application in fish anaesthesia. *Applied Ichthyology*, 28(4), 553–564.
- Wagner, E., Arndt, R. and Hilton, B., 2002.** Physiological stress responses, egg survival and sperm motility for rainbow trout broodstock anesthetized with clove oil, tricaine methanesulfonate or carbon dioxide. *Aquaculture*, 211, 353–366.
- Wagner, G.N., Singer, T.D. and McKinley, R.S., 2003.** The ability of clove oil and MS-222 to minimize handling stress in rainbow trout (*Oncorhynchus mykiss* Walbaum). *Aquaculture Research*, 34, 1139–1146.
- Wagner, T. and Congleton, J.L., 2004.** Blood chemistry correlates of nutritional condition, tissue damage, and stress in migrating juvenile Chinook salmon (*Oncorhynchus tshawytscha*). *Canadian Journal of Fisheries and Aquatic Science*, 61, 1066–1074.
- Wendelaar Bonga, S.E., 1997.** The stress response in fish. *Physiological Reviews*, 7, 591–625.
- Williams, T.D., Christiansen, J. and Nygren, S., 1988.** Intermuscular anesthesia of teleost and elasmobranchs using ketamine hydrochloride. Proc. West. Reg. Am. Assoc. Zool. Pk. Aqu. Monterey Bay Aquarium, Monterey, CA. pp. 132–135.
- Witeska, M., Dudyk, N. and Jarkiewicz, J., 2015.** Haematological effects of 2-phenoxyethanol and etomidate in carp (*Cyprinus carpio* L.). *Veterinary Anaesthesia and Analgesia*, 42, 537–546.