

Research Article

Comparative genotoxic and histopathological effects of copper nanoparticles and copper chloride in goldfish (*Carassius auratus*)Yavaş C.^{1*}, Gülsoy N.²

¹Department of Molecular Biology and Genetics, Faculty of Engineering and Natural Sciences, Biruni University, Istanbul, Türkiye

²Department of Biology, Faculty of Sciences, Marmara University, Istanbul, Türkiye

* Correspondence: cyavas@biruni.edu.tr

Keywords

Copper,
Micronucleus,
Gill histopathology,
Genotoxicity,
Comet assay,
Toxicity

Abstract

The purpose of the current study was to ascertain the comparative effects of compounds, copper (II) oxide nanoparticles (nCuO) and copper salt (CuCl₂), on the genotoxicity and histopathology of goldfish (*Carassius auratus*) for acute exposures of 24, 48, 72, and 96 h (concentrations: 100, 150, 200, and 1, 1.5, 2 mg/L nCuO and CuCl₂, respectively). The peripheral erythrocytes were obtained for that purpose from the caudal vein in each individual. The Comet assay and micronucleus test were used to evaluate genotoxicity, and gill tissues were prepared for histopathology. High levels of nCuO and CuCl₂ concentrations and acute exposure (96-hour) increased the percentage of DNA in the comet tails, olive tail moments, and micronucleus counts showed genotoxic effects on exposed fish. Also, the gill tissue alterations were observed and changed as the lifting of the respiratory epithelium, hyperplasia of the lamellar epithelium, incomplete, and complete fusion of several lamellae, and hypertrophy. Genotoxicity for nCuO and CuCl₂ was discovered to be a concentration and time-dependent phenomenon. In general, experimental groups showed significant impacts ($p=0.05$) on concentrations and exposure periods, which may contribute to understanding the mechanism of nCuO and CuCl₂-induced genotoxicity in the fish.

Article info

Received: January 2024

Accepted: April 2024

Published: May 2024



Copyright: © 2023 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

Introduction

Water pollution is a global problem caused by anthropogenic activities such as industrial discharge, agricultural runoff, and urbanization, which endanger aquatic ecosystems and especially fish populations. Among numerous pollutants such as heavy metals, herbicides, and chemical compounds, oxidative stress has been identified as an important mechanism (Fuat Gulhan *et al.*, 2012; Selamoglu *et al.*, 2015; Caglar *et al.*, 2019). This stress results from an imbalance between reactive oxygen species (ROS) production and antioxidant defenses activated by pollutants through direct ROS generation, enzyme stimulation and antioxidant system disruption. Fish respond to oxidative stress by altering enzyme activity, biomarker levels and gene expression, eliciting physiological adaptations and disease (Orun *et al.*, 2005; Talas *et al.*, 2008; Selamoglu, 2011; Kakoolaki *et al.*, 2013; Talas *et al.*, 2014).

To maintain the long-term viability of aquatic ecosystems, mitigation techniques include environmental management, regulatory policy and multidisciplinary research. Aquatic products and animals are indispensable components of the human diet and play a crucial role in the global aquatic product industry, serving consumers worldwide. Therefore, safeguarding our aquatic environments against pollution is imperative due to its diverse environmental and ecological ramifications. The emissions of volatile organic substances and the contamination of water by oil chemicals and various hazardous agents pose significant threats to aquatic ecosystems and the organisms inhabiting them. Among these organisms,

fish stand out as essential nutrients in the human diet and as a prominent commodity in the global aquatic product market. Consequently, there is an urgent need to prioritize the enhancement of aquatic products and their health. This necessitates a concerted focus on oxidative stress studies and the adoption of sustainable practices aimed at protecting fish populations and preserving the integrity of aquatic ecosystems (Ates *et al.*, 2008; Kakoolaki *et al.*, 2013; Mesut, 2021; Selamoglu, 2021). By proactively addressing these challenges, we can reduce the negative impacts of pollution on aquatic environments while maintaining high quality aquatic products and fish species.

One of the most prevalent transition metals in nature, copper (Cu) serves a variety of functions in organisms and is a necessary nutrient (Burke and Handy, 2005). Cu contamination is pervasive in the aquatic environment. Copper is frequently found in aquatic systems from both natural and man-made sources. Geological deposits, volcanic activity, weathering and erosion of rocks and soils, and geological deposits are all natural sources of copper in surface water. Aquatic system pollution is a serious environmental issue that is becoming more and more global. Aside from cadmium, copper is one of the most often used metals that cause pollution. In several industrial and agricultural operations, people use copper (Carvalho and Fernandes, 2008; Kumar *et al.*, 2021). It is a part of several fungicides, algacides, bactericides, and herbicides, particularly in the forms of sulfate, hydroxide, oxychloride, carbonate, citrate, or gluconate (Ali *et al.*, 2019). Therefore,

contamination will unavoidably happen by Cu forms. Due to human and natural inputs, waterborne Cu has been identified in the environment in high amounts (Cu, 0.04-560 g/L) (Agency, 2007; Oliveira *et al.*, 2008; Yusni and Ifanda, 2020).

Due to their remarkable physicochemical characteristics, metal oxide nanoparticles (NPs) are often employed in a wide range of consumer goods and industrial technologies. Nanomaterials are employed in a variety of home furnishings and appliances, the production of textiles and electronics, the creation of medical devices, and bioremediation technology (Montes *et al.*, 2012). The items incorporating nanomaterials will be released into aquatic ecosystems and agricultural areas as a result of large-scale manufacture and use

(Chatterjee, 2008; Fabrega *et al.*, 2011; Scott-Fordsmand *et al.*, 2017). Metal oxide NPs are among the most popular nanomaterials, and their possible ecological consequences have attracted a lot of interest. Due to their increased potential for potential uses in the future and the fact that NPs are an addition to personal care products, copper (II) oxide nanoparticles (nCuO) are produced in huge quantities for both industrial and domestic usage (Hochella *et al.*, 2019; Ming *et al.*, 2020). Additionally, CuO NPs may find use in a variety of industrial settings (Fig. 1). CuO NPs are thus among the most likely environmental pollutants (Nel *et al.*, 2006; Klaine *et al.*, 2008; Poynton *et al.*, 2011; Hou *et al.*, 2017).




Figure 1: A scanning electron microscope (SEM) image of nCuO.

The significance of determining the toxicological interactions and effects of nCuO and copper salt (CuCl_2) combinations on aquatic life is highlighted by the widespread copper pollution. In this study, goldfish are used as a model to evaluate the toxic effects of the combined

exposure of nCuO and CuCl_2 in freshwater ecosystems on fish. To investigate the potential genotoxic and cytotoxic effects in both structures of copper we used Comet and micronucleus assays as a test for DNA damage and gill tissue for histopathology. The significance of determining the

toxicological interactions and effects of nCuO and CuCl₂ combinations on aquatic life is highlighted by the widespread copper pollution.

Materials and methods

Following the commencement of the experiment, the total copper (Cu) contents of all experimental groups were assessed using an inductively coupled plasma optical emission spectrometer (ICP-OES) (at the 1 hour). The goldfish (*Carassius auratus*) were purchased from a commercial supplier (Dogasan Aquarium, Istanbul, Turkey) and were transferred to the laboratory. They were given the commercial base diet of *C. auratus* (38.7% crude protein; 13% crude fat; 14.8% crude fiber ash) and acclimated for a month at 26±1°C under natural photoperiod. At the start and the conclusion of each exposure session, the standard water quality parameters were tested according to established procedures (APHA, AWWA, WPCF 2005). The pH ranged from 7.6 to 8.1±0.1, dissolved oxygen was at 8.3±0.3 mg/L, nitrite was at 0.3 mg/L, and ammonium was at 0.05 mg/l. In this study, 180 goldfish from both sexes (the mean weight and length of 0.72±0.1 g and 4.7±0.1 cm, respectively) were prepared. The experiments were randomly performed on six groups and at four different times, together with a positive (5 mg/L ethyl methanesulfonate) and negative control in three duplicate tanks. There were 10 fish in each tank (40×30×50 cm) with 25 L of dechlorinated tap water. Fish were not fed during the experiments in order not to cause physiological changes.

nCuO (Alfa Aesar, K02S018, Germany) and CuCl₂ (Alfa Aesar, 10154919,

Germany) were produced in diluted water and supplied to the aquariums. Specific particle size distribution of nCuO nanometallic particles with scanning determined by electron micrographs (SEM). Subnanometer-sized nanoparticles with solubility, as determined from SEM images, ensure no aggregation is observed. It has been found to be about 30nm in the regions. This result is commercially obtained nano copper (II) is compatible with the property given in the material data sheet of the oxide (Fig. 1). For the acute studies, four distinct nCuO and CuCl₂ concentrations (nominal concentrations: 100, 150, 200, and 1, 1.5, 2 mg/L for nCuO and CuCl₂) and four time periods (24, 48, 72, and 96 h). The acute values of Copper compounds in goldfish, presented as 96-h median lethal doses, were used to choose the test concentrations (Kahru and Dubourguier, 2010).

To reduce changes brought on by fish metabolism, the volatilization of less stable chemicals, and organism catabolites, the test water was replaced every 24 hours (80%) by providing semi-static test conditions. After exposure times, blood samples were taken from the caudal vena of anesthetized fish using heparinized syringes (100 mg/L MS-222, Sigma Aldrich). The fish were put back in their tank after a five-minute recovery time in well water.

Genotoxicity

Trypan blue dye was used to examine the cells' viability before the experiment began. Goldfish were used as test subjects for the comet assay, which measures DNA strand breaks (single-strand breaks and alkali-

labile sites). Two fish were removed from each aquarium at each sample period after 24, 48, 72, and 96 hours of exposure and given buffered MS-222 anesthesia. To avoid causing more DNA damage, all of the experimental procedures were carried out in the dark beneath a yellow bulb. The alkaline comet test was carried out using a modified version of the Tice *et al.* Technique (Tice *et al.*, 2000). Briefly, the erythrocytes (10 μ L) were resuspended in cooled PBS (pH 7.4) buffer after the peripheral blood samples were taken from the caudal vasculature using a 1 mL heparinized syringe. The suspension (65 μ L) was then combined with 100 μ L of 0.65% (w/v) agarose with a normal melting point before being applied on a frosted slide that had already been precoated with 0.65% (w/v) agarose with a high melting point. The microscope slide was then submerged in a cold (4°C) lysate solution that contained recently added 1% Triton X100 and 10% DMSO and contained 2.5 M NaCl, 10 mM Na₂EDTA, 10 mM Tris-HCl, and 1% SDS. To facilitate DNA unwinding, the slide was incubated in a newly made alkaline buffer (1 mM Na₂EDTA and 300 mM NaOH, pH 13) for 30 min after 2 h. In the same buffer, electrophoresis was performed at 4 °C for 30 min at 15 V and 300 mA. After being neutralized with a 0.4 M Tris solution (pH 7.59), the slides were immediately stained with 75 L EtBr (10 g/mL) for viewing under a fluorescence microscope (BX51TF, Olympus, Japan) with a 530 nm excitation filter, a 590 nm emission filter, a digital camera (Kameram A640 FL). A computer-based image processing system was used to make the genotoxic observations and analysis

(Kameram Comet Module, Micro System Ltd. Turkey). For data analysis of DNA damage percentage in the tail and olive tail moment, 100 cells from each replication slide were randomly chosen. The fraction of DNA-damaged cells with more than 5% damaged DNA in the tail was used to establish the DNA damage percentage. The amount of DNA in the tail distribution and the distance between the head and the tail were calculated to create the Olive tail moment (Olive *et al.*, 2012). To reduce score fluctuation, each slide was examined by a single observer while remaining completely blind. All chemicals used in these processes were bought from Sigma-Aldrich.

Micronucleus

The slides that would be stained with fish blood were maintained in a solution of 99% ethanol and hydrochloric acid for 10 minutes before being rinsed with distilled water (dH₂O). The slides' edges were covered with fish blood samples. After soaking in 99% ethyl alcohol for ten minutes, the examples were fixed. The slides were run through dH₂O after the ready-made 5% Giemsa (Sigma-Aldrich) stain was applied to the smear preparations. Canadian balsam (Entellan) was used to seal the stained preparations so they could be examined under a light microscope (BX51TF, Olympus, Japan). Four distinct zones were chosen and 250 cells were counted with the 100 x lens. The images were captured using the Image Pro Express 6.0 software. To quantify the frequency of notched nuclei, lobed nuclei, budding, fragmenting, and micronucleated (MN) cells per 1000 cells (%), 1000 erythrocytes

were scored from each slide at a magnification of 1000 x. %MN is determined by dividing the total number of cells by the number of cells that contain MN.

Histopathology

After 96 hours, fish from the negative control group (negative) and nCuO and CuCl₂ exposure groups were removed and dissected gill tissues were processed for histological analysis. Fixation was made in 10% neutral formalin with 0.03% eosin (F5304, Sigma) 24. After fixation for 24 h, gill tissues were dehydrated through a graded series of ethanol for 30 minutes each, cleared in xylene, and infiltrated in paraffin (56-58°C). Sections of 7-10 µm were prepared from paraffin blocks by using a rotary microtome. Staining with Hematoxylin dye (Hematoxylin Solution, Mayer's) for 15 minutes and in Eosin solution (EosinYY-Solution 0.5% Aqueous for Microscopy, Merck) for 5 minutes. Histological preparations were randomly examined three times, and the results from each observation were combined for the final results. Histopathological images were captured using an Olympus BH2 microscope with an image analyzer system (Media Cybernetics, Silver Spring, MD).

Statistical Analysis

The averages and standard deviations (SD) of each independent experiment that was carried out in triplicate were given as experimental data. SPSS for Windows version 11.0 was used to conduct all statistical analyses (SPSS Inc., Chicago, IL, USA). The difference between groups was examined using a three-independent-

samples test, which was followed by a non-parametric Mann-Whitney U test. *p*-value 0.05, significance was determined.

Results

In the exposure groups (nCuO: 100, 150, and 200 mg/L and CuCl₂: 1, 1.5, and 2 mg/L) of goldfish at 24, 48, 72, and 96 h with positive, negative controls, comet formations reflecting the DNA profile migrated out of the nucleus was shown by fluorescence staining. When the comet images obtained from the positive control and application groups were compared with the negative control, it was seen that the tail formation increased significantly (*p*=0.05). The Comet images were analyzed using a computer program, and DNA damage levels were presented as % DNA tail and olive tail moment. We discovered that the specimens that were exposed to all concentrations of nCuO and CuCl₂ had significantly more DNA damage than the negative and positive control after performing the comet assay on erythrocytes (Fig. 2). We discovered high comet scores when we compared the positive control (66 ±5.43% Tail DNA and 46±3.21 Olive tail moment) and experimental groups. The highest comet score compared to the positive control was seen in fish exposed to nCuO. It was observed that the experimental groups exposed to both chemicals had higher DNA damage than the positive control (Table 1, Figs. 2 to 4)

After staining and monitoring under the light microscope, the frequencies of MNs generated outside the main nucleus as a result of genetic damage were assessed. When the MN pictures from the positive, negative, and experimental groups were

compared to the negative and positive controls, MN formation was determined to be significantly increased ($p=0.05$). The quantity of MN increased rapidly, especially as the exposure period increased (Fig. 5).

Histological images of the gills obtained from the positive, negative, and experimental groups were compared with the negative and positive control.

Table 1: nCuO and CuCl₂ at various concentrations have an impact on comet properties in the erythrocytes of *Carassius auratus*.

	Concentration	Time	Tail % DNA	Olive tail moment
	Negative		28 ± 1.3	10 ± 0.7
	Positive	24h	66 ± 5.43	46 ± 3.21
nCuO	100 mg/L	24h	65 ± 3.78	52 ± 1.36
		48h	74 ± 3.9	59 ± 2.11
		72h	78 ± 3.45	66 ± 2.21
		96h	81 ± 4.45	73 ± 3.3
	150 mg/L	24h	71 ± 3.65	59 ± 1.7
		48h	79 ± 2.7	80 ± 4.87
		72h	77 ± 3.9	66 ± 2.4
		96h	85 ± 3.15	76 ± 2.6
	200 mg/L	24h	69 ± 2.21	58 ± 1.91
		48h	73 ± 4.6	65 ± 4.21
		72h	80 ± 3.57	68 ± 2.13
		96h	86 ± 3.15	79 ± 3.78
1 mg/L	24h	76 ± 2.78	52 ± 2.9	
	48h	61 ± 2.96	59 ± 2.87	
	72h	81 ± 5.12	66 ± 2.45	
	96h	80 ± 3.38	73 ± 2.24	
CuCl ₂	1.5 mg/L	24h	78 ± 2.7	59 ± 1.36
		48h	62 ± 2.6	80 ± 3.91
		72h	74 ± 2.13	66 ± 2.74
		96h	82 ± 4.78	76 ± 2.87
	2 mg/L	24h	76 ± 2.9	58 ± 1.5
		48h	65 ± 2.78	65 ± 2.04
		72h	82 ± 3.46	68 ± 2.19
		96h	80 ± 5.24	79 ± 3.21




Figure 2: Erythrocytes of *C. auratus* showing: (a) negative control DNA, (b) positive control DNA, (c) DNA damaged after exposure to nCuO and (d) DNA damaged after exposure to CuCl₂.




Figure 3: Showing tail % DNA and standard error bars as a result of exposure to nCuO at different concentrations and durations with positive and negative control in goldfish ($p < 0.05$).




Figure 4: Showing tail % DNA and standard error bars as a result of exposure to CuCl₂ at different concentrations and durations with positive and negative control in goldfish ($p < 0.05$).




Figure 5: MN counts are shown on a graph of comparative frequency in the erythrocytes of goldfish after exposure to nCuO and CuCl₂ ($p < 0.05$).

Significant deteriorations in the gill epithelium such as uplift, edema, fusion of secondary lamellae, hyperplasia, and hypertrophy were observed that would

prevent even the function of the gills (Table 2). The images obtained as a result of histopathological staining revealed the differentiation in the gill structure (Fig. 6).

Table 2: Histopathological evaluation of the findings of nCuO and CuCl₂ administration groups.

Concentration	nCuO			CuCl ₂		
	100 mg/L	150 mg/L	200 mg/L	1 mg/L	1.5 mg/L	2mg/L
Degeneration of secondary lamellae	+	+	++	+	+	+
Epithelial lift	+	++	+++	+	++	++
Fusion	+	++	+++	+	++	++
Hyperplasia	+	++	++	+	+	++
Hypertrophy	+	++	+++	++	+	++
Edema	-	+	++	-	+	+

+:less, ++:moderate, +++:severe




Figure 6: Micrograph showing the gill histopathology of goldfish treated with nCuO and CuCl₂ respectively a) mucous hypertrophy b) lamella epithelial removal c) fusion of four secondary lamellae.

Discussion

The primary objective of this study was to systematically collect and analyze data to elucidate the genotoxic and histopathological impacts of nCuO and CuCl₂ on *Carassius auratus*, commonly known as goldfish. This experiment aimed to determine the differential effects of these copper-based compounds at various concentrations, exploring their potential to induce DNA damage and cellular alterations within the tissues of the exposed organisms. By employing a range of biomarkers for genotoxicity and histopathological assessment, the study sought to provide a comprehensive overview of the cellular and molecular disruptions occurring in response to these specific contaminants. The results of this research are intended to contribute valuable insights into the toxicological profiles of nCuO and CuCl₂, thereby informing risk assessment and management strategies in aquatic environments where these compounds are prevalent contaminants (Kahru and Dubourguier, 2010).

Behavioral changes in fish were monitored throughout chemical administration (Sharma *et al.*, 2019). In this study, it was found that since there was an excessively dark environment in all concentrations of nCuO, the fish swam close to the surface of the water and there was a decrease in their movement. In CuCl₂ concentrations, active swimming rates decreased and they moved more towards the bottom. In a similar study, Tilton *et al.* reported that active swimming rates of zebrafish, in a mixture of copper chloride and chlorpyrifos, decreased as in their study (Tilton *et al.*, 2011).

Results showed that the comet test was more sensitive in fish studies than the micronucleus test, and they suggested that the two tests, which revealed DNA damage from distinct perspectives, may be combined (Ternjej *et al.*, 2010; Pavlica *et al.*, 2011). The integrated biomarker response index, which integrates the numerous biomarkers in a multivariate data set, can offer a comprehensive harmful effect of pollution on organisms. Many

field and laboratory environmental risk assessment studies have used this strategy. However, till recently, no investigations on the harmful effects of TiO₂ NPs at projected ecologically relevant concentrations on marine scallop have been published (Kim *et al.*, 2010; Kim *et al.*, 2013; Kim and Jung, 2016). Blood cells are susceptible to potentially harmful compounds, as indicated by research involving CuSO₄ molecules (Beninca *et al.*, 2012). Researchers who performed the comet test to demonstrate genotoxicity discovered time- and concentration-dependent increases in the exposed groups as compared to the control groups by examining the proportion of DNA tail damage (Ghisi Nde *et al.*, 2011; Pavlica *et al.*, 2011; Beninca *et al.*, 2012). Many researchers have shown that the comet test is an inexpensive, rapid and reliable test for detecting genotoxicity (Mohanty *et al.*, 2011; Sponchiado *et al.*, 2011; Kousar and Javed, 2015; Chelomin *et al.*, 2017; Kaygisiz and Ciğerci, 2017; Sehirli *et al.*, 2017; Jafar *et al.*, 2019; Atila *et al.*, 2020; Moller *et al.*, 2020; Boyadzhiev *et al.*, 2022). In our study, as a result of the examinations made with the comet test, DNA damages were more clearly seen with the increase in concentration and exposure time, in accordance with the literature. Our results showed that there was more damage in the exposed groups compared to the positive control. Studies have shown that copper nanomaterials cause more DNA damage compared to other nanomaterials. It has also been found that the comet test is sensitive and accurate in revealing potential nanomaterial DNA damage (Boyadzhiev *et al.*, 2022). Studies have shown that copper

nanomaterials cause more DNA damage compared to other nanomaterials. They have also found that the comet test is sensitive and accurate in revealing potential nanomaterial DNA damage (Boyadzhiev *et al.*, 2022).

The studies demonstrated how a variety of heavy metals increased MN counts and can be used on blood and other tissue cells (Bolognesi and Hayashi, 2011; Obiakor *et al.*, 2012; Di Bucchianico *et al.*, 2013). Depending on the concentration and duration used during the application of the nCuO and CuCl₂ linear increase in MN counts was found (Fig. 5). This increase was observed to be greater in fish exposed to nCuO. Semisch *et al.* explained that nCuO particles accumulate more in the nucleus than other copper forms, resulting in an increased level of DNA strand breaks (Semisch *et al.*, 2014). A Genotoxic study of *Labeo rohita* revealed the development of micronuclei, supporting the MN test as a biomarker tool for water contamination (Hussain *et al.*, 2018). It is worth mentioning that the formation of these nuclear anomalies caused by toxicant exposures may further require explainable studies (Cavas and Ergene-Gozukara, 2003).

Gill tissues are significant target tissues for histopathological alterations because they play a fundamental role in respiration and are in direct contact with water (Wood and Eom, 2021). Copper exposure is known to cause alterations in the gill epithelium such as hypertrophy, hyperplasia, necrosis, and secondary lamellae fusion (Karayakar *et al.*, 2010; Ostaszewska *et al.*, 2016). Furthermore, copper nanoparticles build up more quickly in the liver and other organs

of living things than in bone tissues (Wang *et al.*, 2014). Not only copper forms do not damage tissues and organs, but studies have also reported that pesticides used in agriculture also damage different tissues such as the brain. Continuous use of pesticides enters the aquatic ecosystem and threatens the living organisms in the water. According to current understanding, living organisms may experience oxidative stress as a consequence of the accumulation of hazardous chemicals (Ostaszewska *et al.*, 2016). In this case, the air supply will decrease, it will disrupt organ functions, shorten the lifespan of the living thing significantly, and even cause death. (Luzio *et al.*, 2013). Zebrafish gill alterations revealed that epithelial cell proliferation as well as edema of main and secondary gill filaments following exposure after 48 hours of exposure to 0.25 mg/L ionic copper and 1.5 mg/L Cu NPs in the literature (Griffitt *et al.*, 2007; Griffitt *et al.*, 2008; Griffitt *et al.*, 2009). In our study, we observed degeneration of secondary lamellae, epithelial lift, fusions, hyperplasia, hypertrophy, and edema. These alterations contribute to thickening and damaging the surface of the gills which inhibates gas exchange, particularly oxygen uptake. (Al-Bairuty *et al.*, 2013; Hao *et al.*, 2013). As a result of our study, tissue alterations and damages seen in fish exposed to nCuO may have a greater impact on the overall health of aquatic organisms than CuCl₂.

Given the critical role of essential metals (Fe, Zn, and Cu) as cofactors in numerous enzymatic processes, their levels in fish are significant due to strict physiological regulation, emphasizing the importance of understanding these concentrations for both

fish management and human consumption (Kamaruzzaman *et al.*, 2011). They have been reported that the accumulation of these metals at certain levels is not fatal for living organisms or when they are transferred through the food chain. However, the damage that will occur in the accumulation of metals in living organisms can reach levels that can no longer be tolerated by living organisms. These accumulations are seen in different tissues (Caglar *et al.*, 2019). Metal accumulation in fish differs between tissues and is affected by the environmental conditions of the environment where the fish live (Unlu *et al.*, 2009). Heavy metals have been found to be more dangerous for aquatic ecosystems, especially for fish feeding in deep waters (Popek *et al.*, 2003). After heavy metal accumulation in fish, particularly through the food chain, toxic effects ensue in other organisms. Nonetheless, studies have clearly indicated that the detrimental effects of heavy metal accumulation in fish can be prevented by certain antioxidative substances. One of the most frequently reported antioxidative substances is propolis. (Fuat Gulhan *et al.*, 2012; Kakoolaki *et al.*, 2013; Talas *et al.*, 2014; Selamoglu *et al.*, 2015) and selenium (Ates *et al.*, 2008; Talas *et al.*, 2008; Selamoglu, 2011).

Exposure of goldfish to nCuO and CuCl₂ at relevant concentrations resulted in DNA damages as determined by comet assay of genotoxicity. Histopathological examination revealed that the gill is a target organ for copper toxicity. nCuO caused more gill morphological changes upon exposure, than CuCl₂. Both types of copper forms can produce increased oxidative

stress, leading to DNA damage in the case of exposure to copper. MN test is as sensitive as the comet test in detecting genotoxicity. However, more techniques should be used to quantify genotoxic capacity. As a result, copper oxide nanoparticles (nCuO) and copper salt (CuCl₂) can cause harmful effects in fish at these concentrations. The release of excess copper concentration into the aquatic environment poses a potential risk to the aquatic environment. As a result of copper accumulation in aquatic organisms, DNA and tissue damage caused by excess copper accumulation, which they transferred to humans through the food chain, can be reduced by propolis, selenium, and different protective compounds reported in the literature.

Acknowledgments

This publication is derived from the master thesis of Cuneyd Yavas and Marmara University Scientific Research Project Coordination Unit (BAPKO) supported this study, Project No. Fen-A-060510-0139.

Conflict of interest

The authors declare that they have no conflicts of interest.

References

- Agency, U.E.P., 2007.** *Aquatic life criteria -copper.* Available at: <https://www.epa.gov/wqc/aquatic-life-criteria-copper> (Retrieved 10.08.2022)
- Al-Bairuty, G.A., Shaw, B.J., Handy, R.D. and Henry, T.B., 2013.** Histopathological effects of waterborne copper nanoparticles and copper sulphate on the organs of rainbow trout (*Oncorhynchus mykiss*). *Aquatic Toxicology*, 126, 104-115. DOI:10.1016/j.aquatox.2012.10.005
- Ali, H., Khan, E. and Ilahi, I., 2019.** Environmental chemistry and ecotoxicology of hazardous heavy metals: environmental persistence, toxicity, and bioaccumulation. *Journal of Chemistry*, 1-14. DOI:10.1155/2019/6730305
- Ates, B., Orun, I., Talas, Z.S., Durmaz, G. and Yilmaz, I., 2008.** Effects of sodium selenite on some biochemical and hematological parameters of rainbow trout (*Oncorhynchus mykiss* Walbaum, 1792) exposed to Pb²⁺ and Cu²⁺. *Fish Physiology and Biochemistry*, 34, 53-59. DOI:10.1007/s10695-007-9146-5
- Atila, U., Baydilli, Y.Y., Sehirli, E. and Turan, M.K., 2020.** Classification of DNA damages on segmented comet assay images using convolutional neural network. *Computer Methods and Programs in Biomedicine*, 186, 105-192. DOI:10.1016/j.cmpb.2019.105192
- Beninca, C., Ramsdorf, W., Vicari, T., de Oliveira Ribeiro, C.A., de Almeida, M.I., Silva de Assis, H.C. and Cestari, M.M., 2012.** Chronic genetic damages in *Geophagus brasiliensis* exposed to anthropic impact in estuarine lakes at Santa Catarina coast--southern of Brazil. *Environmental Monitoring and Assessment*, 184, 2045-2056. DOI:10.1007/s10661-011-2098-3
- Bolognesi, C. and Hayashi, M., 2011.** Micronucleus assay in aquatic animals. *Mutagenesis*, 26, 205-213. DOI:10.1093/mutage/geq073

- Boyadzhiev, A., Solorio-Rodriguez, S.A., Wu, D., Avramescu, M.L., Rasmussen, P. and Halappanavar, S., 2022.** The High-Throughput In Vitro CometChip Assay for the Analysis of Metal Oxide Nanomaterial Induced DNA Damage. *Nanomaterials (Basel)*, 12, 18-44. DOI:10.3390/nano12111844
- Burke, J. and Handy, R.D., 2005.** Sodium-sensitive and -insensitive copper accumulation by isolated intestinal cells of rainbow trout *Oncorhynchus mykiss*. *Journal of Experimental Biology*, 208, 391-407. DOI:10.1242/jeb.01379
- Caglar, M., Canpolat, O. and Selamoglu, Z., 2019.** Determination of some heavy metal levels in three freshwater fish in Keban Dam Lake (Turkey) for public consumption. *Iranian Journal of Fisheries Sciences*, 18, 188-198. DOI:10.22092/IJFS.2018.117890
- Carvalho, C.D.S. and Fernandes, M.N., 2008.** Effect of copper on liver key enzymes of anaerobic glucose metabolism from freshwater tropical fish *Prochilodus lineatus*. *Comparative Biochemistry and Physiology Part A: Molecular & Integrative Physiology*, 151, 437-442. DOI:10.1016/j.cbpa.2007.04.016
- Cavas, T., and Ergene-Gozukara, S., 2003.** Micronuclei, nuclear lesions and interphase silver-stained nucleolar organizer regions (AgNORs) as cytogenotoxicity indicators in *Oreochromis niloticus* exposed to textile mill effluent. *Mutation Research*, 538, 81-91. DOI:10.1016/s1383-5718(03)00091-3
- Chatterjee, R., 2008.** The challenge of regulating nanomaterials. *Environmental Science and Technology*, 42, 339-343.
- Chelomin, V.P., Slobodskova, V.V., Zakhartsev, M., and Kukla, S., 2017.** Genotoxic potential of copper oxide nanoparticles in the bivalve mollusk *Mytilus trossulus*. *Journal of Ocean University of China*, 16, 339-345. DOI:10.1007/s11802-017-3133-y
- Di Bucchianico, S., Fabbrizi, M.R., Misra, S.K., Valsami-Jones, E., Berhanu, D., Reip, P., Bergamaschi, E., and Migliore, L., 2013.** Multiple cytotoxic and genotoxic effects induced in vitro by differently shaped copper oxide nanomaterials. *Mutagenesis*, 28, 287-299. DOI:10.1093/mutage/get014
- Fabrega, J., Luoma, S.N., Tyler, C.R., Galloway, T.S., and Lead, J.R., 2011.** Silver nanoparticles: behaviour and effects in the aquatic environment. *Environment International*, 37, 517-531. DOI:10.1016/j.envint.2010.10.012
- Fuat Gulhan, M., Duran, A., Selamoglu Talas, Z., Kakoolaki, S., and Mansouri, S., 2012.** Effects of propolis on microbiologic and biochemical parameters of rainbow trout (*Oncorhynchus mykiss*) after exposure to the pesticide. *Iranian Journal of Fisheries Sciences*, 11, 490-503. DOR:20.1001.1.15622916.2012.11.3.4. 4
- Ghisi Nde, C., Ramsdorf, W.A., Ferraro, M.V., de Almeida, M.I., Ribeiro, C.A., and Cestari, M.M., 2011.** Evaluation of genotoxicity in *Rhamdia quelen* (*Pisces, Siluriformes*) after sub-chronic contamination with Fipronil. *Environmental Monitoring and*

- Assessment*, 180, 589-599.
DOI:10.1007/s10661-010-1807-7
- Griffitt, R.J., Weil, R., Hyndman, K.A., Denslow, N.D., Powers, K., Taylor, D., and Barber, D.S., 2007.** Exposure to copper nanoparticles causes gill injury and acute lethality in zebrafish (*Danio rerio*). *Environmental Science and Technology*, 41, 8178-8186.
DOI:10.1021/es071235e
- Griffitt, R.J., Luo, J., Gao, J., Bonzongo, J.C., and Barber, D.S., 2008.** Effects of particle composition and species on toxicity of metallic nanomaterials in aquatic organisms. *Environmental Toxicology and Chemistry: An International Journal*, 27, 1972-1978.
DOI:10.1897/08-002.1
- Griffitt, R.J., Hyndman, K., Denslow, N.D., and Barber, D.S., 2009.** Comparison of molecular and histological changes in zebrafish gills exposed to metallic nanoparticles. *Toxicological Sciences*, 107, 404-415.
DOI:10.1093/toxsci/kfn256
- Hao, L., Chen, L., Hao, J., and Zhong, N., 2013.** Bioaccumulation and sub-acute toxicity of zinc oxide nanoparticles in juvenile carp (*Cyprinus carpio*): a comparative study with its bulk counterparts. *Ecotoxicology and Environmental Safety*, 91, 52-60.
DOI:10.1016/j.ecoenv.2013.01.007
- Hochella, M.F., Jr., Mogk, D.W., Ranville, J., Allen, I.C., Luther, G.W., Marr, L.C., McGrail, B.P., Murayama, M., Qafoku, N.P., Rosso, K.M., Sahai, N., Schroeder, P.A., Vikesland, P., Westerhoff, P., and Yang, Y., 2019.** Natural, incidental, and engineered nanomaterials and their impacts on the Earth system. *Science*, 363, 82-99.
DOI:10.1126/science.aau8299
- Hou, J., Wang, X., Hayat, T., and Wang, X., 2017.** Ecotoxicological effects and mechanism of CuO nanoparticles to individual organisms. *Environmental Pollution*, 221, 209-217.
DOI:10.1016/j.envpol.2016.11.066
- Hussain, B., Sultana, T., Sultana, S., Masoud, M.S., Ahmed, Z., and Mahboob, S., 2018.** Fish ecogenotoxicology: Comet and micronucleus assay in fish erythrocytes as in situ biomarker of freshwater pollution. *Saudi Journal of Biological Sciences*, 25, 393-398.
DOI:10.1016/j.sjbs.2017.11.048
- Jafar, J.R., Attarde, S.S., and Pandit, S.V., 2019.** Individual and combined exposure of cupric chloride di-hydrate and anthracene to *Lamellidens corrianus* (Lea): Evaluation of genotoxic responses using comet assay. *Indian Journal of Experimental Biology*, 57, 949-955.
- Kahru, A., and Dubourguier, H.C., 2010.** From ecotoxicology to nanoecotoxicology. *Toxicology*, 269, 105-119.
DOI: 10.1016/j.tox.2009.08.016
- Kakoolaki, S., Talas, Z.S., Cakir, O., Ciftci, O., and Ozdemir, I., 2013.** Role of propolis on oxidative stress in fish brain. *Basic and Clinical Neuroscience*, 4, 153-158.
- Kamaruzzaman, B., Rina, Z., John, B.A., and Jalal, K., 2011.** Heavy metal accumulation in commercially important fishes of South West Malaysian coast. *Research Journal of Environmental*

- Sciences*, 5, 595-602.
DOI:10.3923/rjes.2011.
- Karayakar, F., Cıçık, B., Ciftci, N., Karaytug, S., Erdem, C., and Ozcan, A., 2010.** Accumulation of copper in liver, gill and muscle tissues of *Anguilla anguilla* (Linnaeus, 1758). *Journal of Animal and Veterinary Advances*, 9, 2271-2274. DOI: 10.3923/javaa.2010.2271.2274
- Kaygisiz, Ş.Y., and Cığerci, İ.H., 2017.** Genotoxic evaluation of different sizes of iron oxide nanoparticles and ionic form by SMART, Allium and comet assay. *Toxicology and Industrial Health*, 33, 802-809. DOI:10.1177/0748233717722907
- Kim, W.K., Lee, S.K., and Jung, J., 2010.** Integrated assessment of biomarker responses in common carp (*Cyprinus carpio*) exposed to perfluorinated organic compounds. *Journal of Hazardous Materials*, 180, 395-400. DOI:10.1016/j.jhazmat.2010.04.044
- Kim, W.K., Lee, S.K., Choi, K., and Jung, J., 2013.** Integrative assessment of biomarker responses in pale chub (*Zacco platypus*) exposed to copper and benzo[a]pyrene. *Ecotoxicology and Environmental Safety*, 92, 71-78. DOI:10.1016/j.ecoenv.2013.02.010
- Kim, W.K., and Jung, J., 2016.** In situ impact assessment of wastewater effluents by integrating multi-level biomarker responses in the pale chub (*Zacco platypus*). *Ecotoxicology and Environmental Safety*, 128, 246-251. DOI:10.1016/j.ecoenv.2016.02.028
- Klaine, S.J., Alvarez, P.J., Batley, G.E., Fernandes, T.F., Handy, R.D., Lyon, D.Y., Mahendra, S., McLaughlin, M.J., and Lead, J.R., 2008.** Nanomaterials in the environment: behavior, fate, bioavailability, and effects. *Environmental Toxicology and Chemistry*, 27, 1825-1851. DOI:10.1897/08-090.1
- Kousar, S., and Javed, M., 2015.** Diagnosis of metals induced DNA damage in fish using comet assay. *Pakistan Veterinary Journal*, 35, 168-172.
- Kumar, V., Pandita, S., Singh Sidhu, G.P., Sharma, A., Khanna, K., Kaur, P., Bali, A.S., and Setia, R., 2021.** Copper bioavailability, uptake, toxicity and tolerance in plants: A comprehensive review. *Chemosphere*, 262, 127810. DOI:10.1016/j.chemosphere.2020.127810
- Luzio, A., Monteiro, S.M., Fontainhas-Fernandes, A.A., Pinto-Carnide, O., Matos, M., and Coimbra, A.M., 2013.** Copper induced upregulation of apoptosis related genes in zebrafish (*Danio rerio*) gill. *Aquatic Toxicology*, 128-129, 183-189. DOI: 10.1016/j.aquatox.2012.12.018
- Mesut, S., 2021.** Blue economy and blue Ocean strategy. *Journal of Ecology and Natural Resources*, 5, 000263. DOI:10.23880/jenr-16000263
- Ming, S., Liu, X., Zhang, W., Xie, Q., Wu, Y., Chen, L., and Wang, H.Q., 2020.** Eco-friendly and stable silver bismuth disulphide quantum dot solar cells via methyl acetate purification and modified ligand exchange. *Journal of Cleaner Production*, 246, 118966. DOI:10.1016/j.jclepro.2019.118966

- Mohanty, G., Mohanty, J., Nayak, A.K., Mohanty, S., and Dutta, S.K., 2011.** Application of comet assay in the study of DNA damage and recovery in rohu (*Labeo rohita*) fingerlings after an exposure to phorate, an organophosphate pesticide. *Ecotoxicology*, 20, 283-292. DOI:10.1007/s10646-010-0580-2
- Moller, P., Muruzabal, D., Bakuradze, T., Richling, E., Bankoglu, E.E., Stopper, H., Langie, S.A.S., Azqueta, A., Jensen, A., Scavone, F., Giovannelli, L., Wojewodzka, M., Kruszewski, M., Valdiglesias, V., Laffon, B., Costa, C., Costa, S., Teixeira, J.P., Marino, M., Del Bo, C., Riso, P., Shaposhnikov, S., and Collins, A., 2020.** Potassium bromate as positive assay control for the Fpg-modified comet assay. *Mutagenesis*, 35, 341-348. DOI:10.1093/mutage/geaa011
- Montes, M.O., Hanna, S.K., Lenihan, H.S., and Keller, A.A., 2012.** Uptake, accumulation, and biotransformation of metal oxide nanoparticles by a marine suspension-feeder. *Journal of Hazardous Materials*, 225-226, 139-145. DOI:10.1016/j.jhazmat.2012.05.009
- Nel, A., Xia, T., Madler, L., and Li, N., 2006.** Toxic potential of materials at the nanolevel. *Science*, 311, 622-627. DOI: 10.1126/science.1114397
- Obiakor, M., Okonkwo, J., Nnabude, P., and Ezeonyejaku, C., 2012.** Eco-genotoxicology: micronucleus assay in fish erythrocytes as in situ aquatic pollution biomarker: a review. *Journal of Animal Science Advances*, 2, 123-133.
- Olive, P.L., Banath, J.P., and Durand, R.E., 2012.** Heterogeneity in radiation-induced DNA damage and repair in tumor and normal cells measured using the "comet" assay. 1990. *Radiation Research*, 178, AV35-42. DOI:10.1667/rrav04.1
- Oliveira, M., Serafim, A., Bebianno, M.J., Pacheco, M., and Santos, M.A., 2008.** European eel (*Anguilla anguilla* L.) metallothionein, endocrine, metabolic and genotoxic responses to copper exposure. *Ecotoxicology and Environmental Safety*, 70, 20-26. DOI:10.1016/j.ecoenv.2007.10.034
- Orun, I., Ates, B., Selamoglu, Z., Yazlak, H., Ozturk, E., and Yilmaz, I., 2005.** Effects of various sodium selenite concentrations on some biochemical and haematological parameters of rainbow trout (*Oncorhynchus mykiss*). *Fresenius Environmental Bulletin*, 14, 18-22.
- Ostaszewska, T., Chojnacki, M., Kamaszewski, M., and Sawosz-Chwalibog, E., 2016.** Histopathological effects of silver and copper nanoparticles on the epidermis, gills, and liver of Siberian sturgeon. *Environmental Science and Pollution Research*, 23, 1621-1633. DOI:10.1007/s11356-015-5391-9
- Pavlica, M., Stambuk, A., Malovic, L., Mladinic, M., and Klobucar, G.I., 2011.** DNA integrity of chub erythrocytes (*Squalius cephalus* L.) as an indicator of pollution-related genotoxicity in the River Sava. *Environmental Monitoring and Assessment*, 177, 85-94. DOI:10.1007/s10661-010-1620-3

- Popek, W., Rosciszewska, M., Biorowiec, F., and Drag-Kozak, E., 2003.** Influence of zinc and copper on development and maturation of gold fish ovaries. *Annals of Animal Science*, 17, 683-687.
- Poynton, H.C., Lazorchak, J.M., Impellitteri, C.A., Smith, M.E., Rogers, K., Patra M., Hammer, K.A., Allen, H.J., and Vulpe, C.D., 2011.** Differential gene expression in *Daphnia magna* suggests distinct modes of action and bioavailability for ZnO nanoparticles and Zn ions. *Environmental Science & Technology*, 45, 762-768. DOI:10.1021/es102501z
- Scott-Fordsmand, J.J., Peijnenburg, W., Semenzin, E., Nowack, B., Hunt, N., Hristozov, D., Marcomini, A., Irfan, M.A., Jimenez, A.S., Landsiedel, R., Tran, L., Oomen, A.G., Bos, P.M.J., and Hund-Rinke, K., 2017.** Environmental Risk Assessment Strategy for Nanomaterials. *International Journal Environmental Research and Public Health*, 14, 1251. DOI:10.3390/ijerph14101251
- Shirli, E., Turan, M.K., and Demiral, E., 2017.** A randomized automated thresholding method to identify comet objects on comet assay images. *Proceedings of the 3rd International Conference on Communication and Information Processing*, 464-467. DOI:10.1145/3162957.3163044
- Selamoglu, M., 2021.** The effects of the Ports and water transportation on the Aquatic ecosystem. *Open Access Journal of Biogenic Science and Research*, 10, 2021. DOI:10.46718/JBGSR.2021.10.000239
- Selamoglu, Z., 2011.** Orun, I., Talas ZS, Alkan, A. Modulating Effect of Selenium on Gills of Fish Exposed to Heavy Metals. *Fresenius Environmental Bulletin*, 20, 2011.
- Selamoglu, Z., Duran, A., Gulhan, M., and Erdemli, M., 2015.** Effects of propolis on biochemical and microbiological parameters in carp (*Cyprinus carpio*) fillets exposed to arsenic. *Iranian Journal of Fisheries Sciences*, 14, 896-907. DOI:20.1001.1.15622916.2015.14.4.7.0
- Selamoglu, M., 2021.** Importance of the cold chain logistics in the marketing process of aquatic products: An update study. *Journal of Survey in Fisheries Sciences*, 25-29. DOI:10.17762/sfs.v8i1.64
- Semisch, A., Ohle, J., Witt, B., and Hartwig, A., 2014.** Cytotoxicity and genotoxicity of nano - and microparticulate copper oxide: role of solubility and intracellular bioavailability. *Particle and Fibre Toxicology*, 11, 10. DOI:10.1186/1743-8977-11-10
- Sharma, M., Thakur, J., and Verma, S., 2019.** Behavioural responses in effect to chemical stress in fish: A review. *International Journal of Fisheries and Aquatic Studies*, 7, 01-05.
- Sponchiado, G., de Lucena Reynaldo, E.M.F., de Andrade, A.C.B., de Vasconcelos, E.C., Adam, M.L., and de Oliveira, C.M.R., 2011.** Genotoxic effects in erythrocytes of *Oreochromis niloticus* exposed to nanograms-per-liter concentration of 17 β -estradiol (E 2): an assessment using micronucleus test and comet assay. *Water, Air, & Soil*

- Pollution*, 218, 353-360.
DOI:10.1007/s11270-010-0649-9
- Talas, Z.S., Orun, I., Ozdemir, I., Erdogan, K., Alkan, A., and Yilmaz, I., 2008.** Antioxidative role of selenium against the toxic effect of heavy metals (Cd^{+2} , Cr^{+3}) on liver of rainbow trout (*Oncorhynchus mykiss* Walbaum 1792). *Fish Physiology and Biochemistry*, 34, 217-222. DOI:10.1007/s10695-007-9179-9
- Talas, Z.S., Gulhan, M.F., Erdogan, K., and Orun, I., 2014.** Antioxidant effects of propolis on carp *Cyprinus carpio* exposed to arsenic: biochemical and histopathologic findings. *Diseases of Aquatic Organism*, 108, 241-249. DOI:10.3354/dao02714
- Ternjej, I., Mihaljevic, Z., Stankovic, I., Kerovec, M., Sipos, L., Zeljezic, D., and Kopjar, N., 2010.** Estimation of DNA integrity in blood cells of eastern mosquitofish (*Gambusia holbrooki*) inhabiting an aluminium-polluted water environment: an alkaline comet assay study. *Archives of Environmental Contamination and Toxicology*, 59, 182-193. DOI:10.1007/s00244-010-9469-6
- Tice, R.R., Agurell, E., Anderson, D., Burlinson, B., Hartmann, A., Kobayashi, H., Miyamae, Y., Rojas, E., Ryu, J.C., and Sasaki, Y.F., 2000.** Single cell gel/comet assay: guidelines for in vitro and in vivo genetic toxicology testing. *Environmental Molecular Mutagenesis*, 35, 206-221. DOI:10.1002/(sici)1098-2280(2000)35:3<206::aid-em8>3.0.co;2-j
- Tilton, F.A., Bammler, T.K., and Gallagher, E.P., 2011.** Swimming impairment and acetylcholinesterase inhibition in zebrafish exposed to copper or chlorpyrifos separately, or as mixtures. *Comparative Biochemistry and Physiology Part C: Toxicology & Pharmacology*, 153, 9-16. DOI:10.1016/j.cbpc.2010.07.008
- Unlu, E., Karadede-Akin, H., Akturk, M.N., and Yanar, M., 2009.** Bioaccumulation of cadmium by Nile tilapia *Oreochromis niloticus* (L.) in the presence of green algae *Cladophora glomerata* (L.) Kutz. *Fresenius Environmental Bulletin*, 18, 2055-2059.
- Wang, T., Long, X., Cheng, Y., Liu, Z., and Yan, S., 2014.** The potential toxicity of copper nanoparticles and copper sulphate on juvenile *Epinephelus coioides*. *Aquatic Toxicology*, 152, 96-104. DOI:10.1016/j.aquatox.2014.03.023
- Wood, C.M., and Eom, J., 2021.** The osmorepiratory compromise in the fish gill. *Comparative Biochemistry and Physiology Part A: Molecular & Integrative Physiology*, 254, 110895. DOI:10.1016/j.cbpa.2021.110895
- Yusni, E., and Ifanda, D., 2020.** Analysis of heavy metal of copper (Cu) and lead (Pb) at Siombak Lake North Sumatera Province. *IOP Conference Series: Earth and Environmental Science*, 454, 012129. DOI:10.1088/1755-1315/454/1/012129