

Research Article

Molecular verification of *Caulerpa racemosa* (Caulerpceae, Chlorophyta) in the Persian Gulf and Oman Sea (south of Iran)

Salehi Balashahri M.¹, Dehshiri M.M.², Sohrabipour J.^{3*}, Mehregan I.¹

1 Department of Biology, Science and Research Branch, Islamic Azad University, Tehran, Iran

2 Department of Biology, Borujerd Branch, Islamic Azad University, Borujerd, Iran

3 Agriculture and Natural Resources Research and Education Center of Hormozgan, Agriculture Research, Education and Extension Organization (AREEO), Bandar Abbas, Iran

*Correspondence: anahitazdp1400@gmail.com

Keywords

Seaweed,
Caulerpceae,
Gulf of Oman,
Persian Gulf,
tufA gene,
Green algae

Abstract

Caulerpa racemosa is edible marine green algae of the Caulerpceae family. *C. racemosa* is consumed as a sea vegetable in many countries, especially Southeast Asian countries. In recent years, molecular studies caused some changes in the taxonomic positions of the members of the genus *Caulerpa*. It is necessary to accurately identify species of this genus based on modern molecular studies due to the consumption of some species and also the toxic effect of some other species of this genus. Some species of *Caulerpa* have already been reported from the southern coasts of Iran based on preliminary morphological studies. The purpose of this study was to collect and accurately identify the species of the genus *Caulerpa* in the coastal waters of the Persian Gulf and the Gulf of Oman in the southern coastlines of Iran. This study was based on the combination of molecular and morphological characteristics studies of the collected samples of this genus in this area. In the current article, the results obtained about the *Caulerpa racemosa* are presented. Samples of the genus were collected from different coastal areas of the Persian Gulf and seashores of Oman Gulf along the geographical borders of Iran and transported to the laboratory in cool conditions. After cleaning and washing, some of the cleaned samples were dried in silica gel and used for DNA extraction and thereof PCR process based on *tufA* gene. In addition, herbarium samples required for morphological studies were prepared and deposited in the herbarium of Agriculture and Natural Resources Research and Education Center of Hormozgan province. The results of this study relatively solve the confusion in *Caulerpa racemosa-peltata* complexity for Iranian members of the complex. In addition to morphological features, the current molecular investigation also helped to accurately identification of the members of this complex in the Iranian seashores of the Persian Gulf and Gulf of Oman. As a highlighted conclusion, the current research on *Caulerpa* species from the Iranian seashores of the Persian Gulf and Gulf of Oman resulted in resolving two species including *Caulerpa sertularioides* (S. G. Gmelin) Howe forma *farlowii* (Weber-Van Bosse) Børgesen, which was resolved as *Caulerpa selago* (Turner) C. Agardh (Salehi Balashahri *et al.*, 2021) and *Caulerpa racemosa* var. *macrophyssa* also resolved as *Caulerpa racemosa* (Forsskål) J. Agardh.

Article info

Received: February 2023

Accepted: June 2023

Published: September 2024



Copyright: © 2023 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

Introduction

Genus *Caulerpa* as a group of family Caulerpaceae in Bryopsidales order comprises around 213 taxa including 104 accepted species, 39 accepted varieties and 70 accepted forma in algaebase (Guiry and Guiry, 2023). Also, a total of 188 taxa including 67 species, 77 variety and 47 forma, currently listed as synonymy on this site. This list of accepted and synonym names shows the high complexity in taxonomy of this genus. Members of this genus are distributed in a wide range of habitats in tropical and subtropical biogeographical marine regions of the world (Meinesz and Boudouresque, 2000; Fama *et al.*, 2002), Phenotypic plasticity is a main source of taxonomic complexity in most algal groups, which led to many misidentifications and nomenclatural confusions and many misestimating in biodiversity (Saunders, 2008; Leliaert *et al.*, 2009; Tronholm *et al.*, 2010), Because of recent expanding of the members of the genus into the temperate marine areas, considerable researches focused on this genus (Meinesz and Hesse, 1991; Piazzini, 2001). Some species such as *caulerpa taxifolia* and *Caulerpa cylindracea* widely spread in the Mediterranean Sea and caused high damage to fish resources in this marine region (Amade *et al.*, 1999).

Members of the *Caulerpa* genus show a wide variety of habits, their thallus morphologically consists of stolons, rhizoids, and upright fronds that have various forms of branchlets or ramuli (Fritsch, 1965). These morphological characteristics have been used to determine species delimitation by phycologists. Environmentally conditions cause many

phenotypic plasticity, especially in the ramuli, which has led to much confusion, high synonyms, and various taxonomic levels such as subspecies, varieties, forms, and ecads (Peterson, 1972, Ohba and Enomoto, 1987; Coppejans and Prud'homme van Reine, 1992).

Molecular tools as a convenient means have provided valuable devices for researchers to delimit and identify species in taxonomically challenging groups (Verbruggen *et al.*, 2009; Saunders, 2008; Leliaert *et al.*, 2009), These tools, however, have caused new difficulties for taxonomists such as conflicts between morphological and molecular concepts (Leliaert *et al.*, 2009; Sauvage *et al.*, 2013) and increasing diversity of cryptic species based on molecular data (De Clerck *et al.*, 2013).

Recent molecular studies on the *Caulerpa* genus have further added difficulties in the taxonomy of this genus because caused some mismatch between morphological and molecular species concepts in the genus, most evident in the *C. racemosa-peltata* complex (Fama *et al.*, 2002; Sauvage *et al.*, 2013; Belton, 2015), All the authors believe that the matching of previously described names with newly obtained molecular data is an extremely difficult task. This research has deposited many misidentified *Caulerpa* specimens on Genbank, which makes it relatively impossible to do accurate taxonomy of the genus. Belton (2015) tried to resolve this problem in the taxonomy of *Caulerpa* genus in Australian waters and provide a framework to integrate traditional nomenclature and molecular-based taxonomies. By using two gene markers

rbcL and *tufA*, Belton (2016) developed this framework for the most problematic taxons of the genus *C. racemosa* (Forsskål) J. Agardh and *C. peltata* J.V. Lamouroux, known as the *C. racemosa–peltata* complex, which has more than 30 described varieties and forms (Guiry and Guiry, 2023).

C. racemosa is composed of a large cell with a large number of nuclei. Chloroplasts contain free chlorophyll that migrates from one part of the body to other parts, and there is a network of fibrous proteins that help to movement of organs. This species is widely found in temperate and tropical shallow marine areas. In 1926 a new form of algae was reported from Tunisia, possibly migrating from the Red Sea, and later migrated to most of the Mediterranean Sea and became more widespread than the invasive species *Caulerpa taxifolia*, and was known as *C. racemosa* var. *cylindracea* which molecular investigation resolved it as *Caulerpa cylindracea* (Wirawan *et al.*, 2022).

C. racemosa is edible seaweed like *Caulerpa lentillifera* and is widely used as a vegetable in Japan, Fiji, Philippines, and Thailand. It is also eaten by local fishermen in Malaysia and Indonesia. This alga is rich in fiber, protein, minerals (calcium and magnesium), folic acid, ascorbic acid, vitamin A, and vitamin B1, while also being low in fat (4), In addition to the nutritional value, *C. racemosa* also has antibacterial and antioxidant properties (Arfini, 2022; Pangestut *et al.*, 2021).

In the Iranian seashores of the Persian Gulf and Gulf of Oman, 11 species and varieties of algae were reported in two reports (Sohrabipour and Rabiei, 1996, 1999) and

in a checklist published by Sohrabipour and Rabiei (2007), which included *Caulerpa fastigiata* Montagne, *Caulerpa manorensis* Nizamuddin, *Caulerpa mexicana* Sonder ex Kützing, *Caulerpa crassifolia* (C. Agardh) J. Agardh, *Caulerpa peltata* J.V. Lamouroux (= *Caulerpa racemosa* var. *peltata* (Lamouroux) Eubank), *Caulerpa racemosa* var. *macrophysa* (Sonder ex Kützing) W.R. Taylor, *Caulerpa sertularioides* (S.G. Gmelin) Howe forma *farlowii* (Weber-Van Bosse) Børgesen, *Caulerpa sertularioides* (S.G. Gmelin) M. Howe forma *Caulerpa sertularioides*, *Caulerpa taxifolia* (M. Vahl) C. Agardh, *Caulerpa brachypus* Harvey, *Caulerpa scalpelliformis* (Turner) C. Agardh. Based on these published papers, Kokabi and Yousefzadii (2015) reported another checklist including the *Caulerpa* species (Shams and Amini, 2017) provided short descriptions for some species of the genus *Caulerpa* collected from southern seashores of Iran.

Materials and methods

Samples of *Caulerpa* were collected from different intertidal regions of the Persian Gulf and Gulf of Oman at the southern coastlines of Iran (Table 1). The collected samples were cleaned and washed thoroughly with seawater to remove sand and other debris then transferred in cool conditions to the laboratory. In the laboratory after precisely cleaning, the samples were washed with tap water and spread on herbarium sheets and small parts of each sample were dried using silica gel and stored at -20°C . The prepared herbarium sheets were deposited in the algal herbarium of the Agriculture and

Natural Resources Research and Education Center of Hormozgan Province, Bandar Abbas, Iran.

Table 1: The geographical information of collecting sites of the samples.

Location	Y	X
Jask	25.636320 N	57.764019 E
	25.640710 N	57.777129 E
Qeshm island	26.981241 N	56.232606 E
	26.969403 N	56.264558E
Chabahar	25.165417 N	61.501707 E
	25.272164 N	60.735824 E

Molecular analysis

DNA extraction was carried out using the modified CTAB method (Promega, Madison, WI, USA) and QiAgen kit. The PCR amplification of the extracted DNA was carried out using *tufA* gene marker based on the protocols proposed by Kazi *et al.* (2013) or Famà *et al.* (2002), The PCR products were sent to Pioneer Technology Gene Company for sequencing. The obtained sequences were then edited with ChromasPro ver. 1.5 (Technelysium Pty Ltd, Australia), After that the newly obtained sequences were blasted in NCBI and the most similar submitted and published sequences (FASTA format) were extracted and aligned in MEGA 7.0 (Kumar *et al.*, 2016) and neighbor-joining phylogenetic tree were constructed.

Results

Morphology of *Caulerpa racemosa* (Forsskål) J. Agardh

The genus of *Caulerpa* is morphologically distinguished by the siphonous stolon

which was attached to the substrate by siphonous rhizoids and had upright assimilators (or fronds) that usually bear characteristic different shape of branchlets (ramuli). Based on these basic morphological features, the taxonomic definition of this genus member in taxonomic levels of the species and variety are determined by morphological variation in shape and configuration of the ramulies (Balton *et al.*, 2014; Riosmeña-Rodriquez *et al.*, 2014).

The samples of the collected *Caulerpa* specimens which were morphologically identified as *C. racemosa* in the current study characterized by short to moderate long erect branches (up to 4 cm) bearing sparse to dense ramulis with short stalks (up to 6 mm height) which, radially, alternately to irregularly were arranged on the erect branch. The distal parts of the ramuli are mainly discoid or truncate shapes, which are up to 8 mm in diameter. The members of the species grow on dead corals covered with sands or on sandy-muddy bottoms at the middle to low intertidal or subtidal zones. In protected or moderately wave-exposed shores they form entangled mats that attach to the sea bottom with densely branched rhizoids. Those growing on rocky-wave exposed reefs possess short erect branches covered with spherical or globose tips ramulies (Fig. 1).

Molecular study

A total set of 50 sequences was obtained from the samples of *Caulerpa* spp. including four sequences from Iran and two sequences of *Caulerpella ambigua* as outgroups were used for phylogenetic analysis. The aligned set composed of 719

base pairs of *tufA* gene nucleotides, which of 530 characters were constant, 38 variable characters were parsimony-uninformative

and 151 were parsimony-informative characters.



Figure 1: Morphological features of the *Caulerpa racemosa* (a) whole morphology of the collected samples, (b) a detailed image from erect frond shows a discoid distal end of ramuli, (c) a detailed image of erect branches shows ramuli on the herbarium sheet (as *C. racemosa* var. *macrophyta* in Sohrabipour and Rabiei, 2007) and (d) a close up image from discoid ends of ramuli of the samples that morphologically already reported as *C. peltata* (Sohrabipour and Rabiei, 2007 and *C. racemosa*, var. *peltata* in Sohrabipour and Rabiei 1999).

Iranian sequences of *Caulerpa racemosa* in phylogenetic analyses grouped in the same clade together with the submitted sequences in GenBank from USA including Florida (DQ652425, DQ652424), New Caledonia (AJ417947, AJ417956), and also, three sequences from Indonesia (FM956052), the Philippines (KT861496), Adelaide (FM956052). The constructed phylogenetic tree was congruent with the

previous phylogenetic analyses which were reported by Fama *et al.* (2002) and Sauvage *et al.* (2013) (Fig. 2).

Intraspecific divergence within the sequences of this clade ranged between 0.0 to 0.5% that is accordance with 3 nucleotides, while interspecific divergence between this clade and clade contains the other varieties of *Caulerpa racemosa* ranged between 2.8-4.4% which was more

than interspecific divergence between sequences of *Caulerpa racemosa* and *Caulerpa selago* (2.5-2.7%) (Table 2). This range of divergence among the varieties of

the *Caulerpa racemosa* may be led to a distinct species in future investigations.



Figure 2: Phylogenetic tree based on *tufA* gene sequences. Numbers above nodes indicate bootstrap support. Support values (>50 is not shown).

Table 2: Divergence ranges within and between the sequences of *Caulerpa* members from Iran and the sequences of *Caulerpa racemosa* from other regions of the world.

	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22
1. T4 <i>Caulerpa racemosa</i> Iran																						
2. T4 <i>Caulerpa racemosa</i> Iran		0.000																				
3. AJ417947C. <i>racemosa</i> var <i>macrophysa</i>		0.002	0.002																			
4. FM956052 <i>C. racemosa</i> <i>macrophysa</i>		0.002	0.002	0.003																		
5. DQ652425 <i>C. racemosa</i>		0.003	0.003	0.002	0.005																	
6. DQ 652424 <i>C. racemosa</i>		0.003	0.003	0.002	0.005	0.000																
7. KT 861496 <i>C. racemosa</i>		0.000	0.000	0.002	0.002	0.003	0.003															
8. AJ417956 <i>C. racemosa</i> var <i>macrophysa</i>		0.003	0.003	0.002	0.005	0.000	0.000	0.003														
9. T3TUFGO407 <i>C. selago</i> Iran		0.027	0.027	0.025	0.025	0.027	0.027	0.027	0.027													
10. T3TUFGO408 <i>C. selago</i> Iran		0.027	0.027	0.025	0.025	0.027	0.027	0.027	0.027	0.000												
11. AJ417973 <i>C. selago</i> Egypt		0.028	0.028	0.027	0.027	0.028	0.028	0.028	0.028	0.002	0.002											
12. AJ417948 <i>C. racemosa</i> var <i>pelteta</i>		0.028	0.028	0.027	0.027	0.029	0.028	0.028	0.028	0.025	0.025	0.027										
13. AJ417957 <i>C. racemosa</i> var <i>lurbinata</i>		0.030	0.030	0.028	0.028	0.030	0.030	0.030	0.030	0.023	0.023	0.025	0.027									
14. JN817684 <i>C. racemosa</i>		0.030	0.030	0.028	0.028	0.030	0.030	0.030	0.030	0.023	0.023	0.025	0.027	0.000								
15. AJ417950 <i>C. racemosa</i>		0.030	0.030	0.028	0.028	0.030	0.030	0.030	0.030	0.027	0.027	0.028	0.002	0.028	0.028							

Table 2 (continued):

	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22
16. AJ417954 <i>C. racemosa</i> var <i>lamourouxii</i>	0.035	0.035	0.033	0.033	0.035	0.035	0.035	0.035	0.028	0.028	0.030	0.033	0.007	0.007	0.035							
17. AJ417955 <i>C. racemosa</i> var <i>occidentalis</i>	0.035	0.035	0.033	0.033	0.035	0.035	0.035	0.035	0.028	0.028	0.030	0.028	0.012	0.012	0.030	0.018						
18. AJ417949 <i>C. racemosa</i> var <i>peltata</i>	0.032	0.032	0.030	0.030	0.032	0.032	0.032	0.032	0.028	0.028	0.030	0.007	0.030	0.030	0.005	0.037	0.032					
19. FM956048 <i>C. racemosa</i> var <i>cylindracea</i>	0.037	0.037	0.035	0.035	0.037	0.037	0.037	0.037	0.030	0.030	0.032	0.030	0.013	0.013	0.032	0.020	0.002	0.033				
20. KJ957101 <i>C. chemnitzia</i>	0.030	0.030	0.028	0.028	0.030	0.030	0.030	0.030	0.027	0.027	0.028	0.005	0.028	0.028	0.003	0.035	0.030	0.002	0.032			
21. KJ957093 <i>C. chemnitzia</i>	0.032	0.032	0.030	0.030	0.032	0.032	0.032	0.032	0.028	0.028	0.030	0.007	0.030	0.030	0.005	0.037	0.032	0.000	0.033	0.002		
22. JN817671 <i>C. chemnitzia</i>	0.030	0.030	0.028	0.028	0.030	0.030	0.030	0.030	0.027	0.027	0.028	0.005	0.028	0.028	0.003	0.035	0.030	0.002	0.032	0.000	0.002	
23. JQ894932 <i>C. racemosa</i>	0.044	0.044	0.024	0.042	0.044	0.044	0.044	0.044	0.033	0.033	0.035	0.037	0.018	0.018	0.038	0.025	0.013	0.040	0.015	0.038	0.040	0.038

Discussion

The previously reported species of the *Caulerpa* genus from the Persian Gulf and the Gulf of Oman indicated the presence of 11 species of the genus from the southern coastline of Iran (Sohrabipour and Rabiei, 2007, 1996, 1999), In the two checklists published by John and Al-Thahni (2015) and Kokabi and Yusefzaadi (2015), totally 13 species of *Caulerpa* were listed by increasing one species *Caulerpa faridii* Nizamuddin from Chabahar at Iranian seashores of Oman gulf (Kokabi and Yusefzaadi, 2015) and one variety *Caulerpa racemosa* [var. *lamourouxii*] f. *requienii* (John and Al-Thahni, 2015) from Abu Dabi. The results of this study relatively solve the confusion in *Caulerpa racemosa-peltata* complex for Iranian

members of the complex. Environmental conditions caused by unstable morphological features have led to confusing nomenclature in taxonomic levels of varieties and forma, of *Caulerpa racemosa-peltata* complex (Sauvage *et al.*, 2013).

Fama *et al.* (2002) in their first molecular analyses based on *tufA* gene data combined with detailed anatomical studies on pyrenoid size, chloroplast size, number of thylakoids in granum, and also number and length of starch grains, showed that the members of clade of *Caulerpa racemosa* have similar detailed morphological features.

The most important visible primary criteria for morphological distinction of the species are the branching pattern of the

assimilator, ramuli shapes, and configuration. These morphological features, however, can show high variation under environmental conditions changes (Peterson, 1972; Calvert, 1976; Ohba and Enomoto, 1987; Ohba *et al.*, 1992; Riosmena-Rodríguez *et al.*, 2014) can lead to considerable taxonomic confusion (Calvert, 1976), therefore using new molecular tools may resolve this complexity.

Some Iranian specimens of the *Caulerpa*, *Caulerpa sertularioides* (S. G. Gmelin) Howe forma *farlowii* (Weber-Van Bosse) Børgesen, which were already reported from various regions of the Iranian and Arabian shores based on the first report of the taxa in this area (Nizamuddin and Gessner, 1970) was resolved as *Caulerpa selago* (Turner) C. Agardh (Salehi Balashahri *et al.*, 2021) and *Caulerpa racemosa* var. *macrophysa* also resolved as *Caulerpa racemosa* (Forsskål) J. Agardh (current study).

The complexity of *Caulerpa racemosa-peltata* complex has already led to many molecular studies based on molecular data focused on *tufA* gene. So the *tufA* (encoding elongation factor TU) of the chloroplast gene as the most common partial gene, generally has been used for the species distinction in the *Caulerpa* genus by almost algal taxonomist in recent years (Famà *et al.*, 2002; Senerpont Domis *et al.*, 2003; Verlaque *et al.*, 2000, 2003, 2004; Durand *et al.*, 2003; Stam *et al.*, 2006; Kazi *et al.*, 2013; Sauvage *et al.*, 2013; Belton *et al.*, 2014, 2015, 2019; Draisma *et al.*, 2014; Fernández-García *et al.*, 2016; Dumilag *et al.*, 2016).

In the current study, based on the molecular data of *tufA* gene, we approved the presence of *Caulerpa racemosa* (Forsskål) J. Agardh that had already been reported as *Caulerpa racemosa* by Sohrabipour and Rabiei (1999, 2007) and also as *Caulerpa racemosa* var. *macrophysa* (Sohrabipour and Rabiei, 2007) from the Iranian coasts of the Persian Gulf and Gulf of Oman. Phylogenetic analysis of *tufA* gene resolved *Caulerpa racemosa* var. *macrophysa* as *Caulerpa racemosa* (Forsskål) J. Agardh (Fig. 2).

Conspecificity of *Caulerpa racemosa* var. *macrophysa* with *Caulerpa racemosa* was already been approved in the previous studies based on *tufA* gene data analyses which combined with morphological studies (Famà *et al.*, 2000, 2002; Verlaque *et al.*, 2000; Stam *et al.*, 2006; Draisma *et al.*, 2014; Sauvage *et al.*, 2013) and phytochemical investigations (Wirawan *et al.*, 2022). Altamirano *et al.* (2014) in their study based on ITS1-5.8S-ITS2 sequences also showed this conspecificity in *Caulerpa racemosa*.

In the recently published paper (Salehi Balashahri *et al.*, 2021) a new record, *Caulerpa selago* (Turner) C. Agardh, also was reported based on the *tufA* data obtained from the gene information of the collected samples of *Caulerpa* specimens that all have already been reported as *Caulerpa sertularioides* (S. G. Gmelin) Howe forma *farlowii* (Weber-Van Bosse) Børgesen from Iranian and Arabian regions of the Persian Gulf (Sohrabipour and Rabiei, 2007, 1996; Kokabi and Yusefzaadi, 2015; John and Al-Thahni, 2015), which all of them followed Nizamuddin and Gessner (1970).

The results of this study showed that modern molecular studies are good tools for the best classification of seaweed species which help to better program planning for economic exploitation of the seaweed resources or environmental issues in marine macroalgae ecological studies concerning the climate change problems that may affect the species richness in marine ecosystems.

Acknowledgments

The authors consider it necessary to express their sincere gratitude to the esteemed management of the General Department of Environmental Protection of Hormozgan Province and the genetic expert of the center, Ms. Mojdeh Ram, who helped us to carry out and improve the quality of this research. Special thanks to Mr. Khosravi, in the Agricultural and Natural Resources Research Center of Hormozgan province, who have helped us in seaweed samples collecting from different regions of the Persian Gulf and the Oman Sea.

Conflicts of interest

The authors have no conflicts of interest to declare. All co-authors have seen and agree with the contents of the manuscript. We certify that the submission is original work and is not under review at any other publication.

References

Altamirano, M., Andreakis, N., Souza-Egipsy, V., Zanolla, M. and De la Rosa, J., 2014. First record of *Caulerpa cylindracea* (Caulerpaceae, Chlorophyta) in Andalusia (Southern Spain). *Anales*

del Jardín Botánico Madrid, 71(2), 1-8. <https://doi.org/10.3989/ajbm.2381>

Amade, M., Tolay, M., Cottalorda, J.M., Vaugelas, J.D. and Meinesz, A., 1999.

Toxicologic effects of the invasive seaweed *Caulerpa taxifolia* in the mediterranean. 4th Congress of Toxicology in Developing Countries, Antalya, Turkey. pp 6-10.

Arfini, F., 2022. Antioxidant and Antibacterial Activity of Caulerpa Flour (*Caulerpa lentifera*) as Functional Food Raw Material. *Asian Food Science Journal*, 22(11):49-54.

DOI: 10.9734/AFSJ/2022/v21i11596

Belton, G.S., Prud'homme van Reine, W.F.M., Huisman, J.M., Draisma, S.G.A. and Gurge, C.F.D., 2014.

Resolving phenotypic plasticity and species designation in the morphologically challenging *Caulerpa racemosa-peltata* complex (Caulerpaceae, Chlorophyta). *Journal of Phycology*, 50, 32-54. DOI: 10.1111/jpy.12132

Belton, G.S., Huisman, J.M. and Gurgel, C.F.D., 2015. Caulerpaceae. In: Huisman J.M. (ed). *Algae of Australia: Marine benthic algae of north-western Australia*. 1. Green and brown algae. Canberra and Melbourne: ABRS and CSIRO Publishing, Australia. pp. 75–102.

Belton, G.S., Draisma, S.G.A., Prud'homme van Reine, W.F., Huisman J.M. and Gurgel, C.F.D., 2019. Taxonomic reassessment of the diversity of *Caulerpa* (Chlorophyta, Caulerpaceae) from southern Australia based on *tufA* and *rbcL* sequence data. *Phycologia*, 52(1), 6–13. DOI: 10.1080/00318884.2018.1542851

Calvert, H.E., 1976. Culture studies on some Florida species of *Caulerpa*:

- morphological responses to reduced illumination. *British Phycological Journal*, 11, 203–214.
DOI:10.1080/00071617600650481
- Ceccherelli, G. and Piazzini, L., 2001.** Dispersal of *Caulerpa racemosa* fragments in the Mediterranean: lack of detachment time effect on Establishment. *Botanica Marina*, 44, 209–213.
<https://doi.org/10.1515/BOT.2001.027>
- Coppejans, E. and Prud'homme van Reine, W.F., 1992.** Seaweeds of the Snellius-II expedition (E. Indonesia), the genus *Caulerpa* (Chlorophyta-Caulerpales), Bull. Séanc. Académie des Sciences D'outre-Mer, 37, 667–712.
- De Clerck, O., Guiry, M.D., Leliaert, F., Samyn, Y. and Verbruggen, H., 2013.** Algal taxonomy: a road to nowhere? *Journal of Phycology*, 49, 215–25.
DOI: 10.1111/jpy.12020
- Draisma, S.G.A., Prud'homme van Reine, W.F., Sauvage, T., Belton, G., Gurgel, C.F.D., Lim, P.E. and Phang, S.M., 2014.** A Re-Assessment of the infra-generic classification of the genus *Caulerpa* (Caulerpaceae, Chlorophyta) inferred from a time-calibrated molecular phylogeny. *Journal of Phycology*, 50, 1020–1034.
<https://doi.org/10.1111/jpy.12231>
- Enomoto, S. and Ohba, H., 1987.** Culture studies on *Caulerpa* (Caulerpales, Chlorophyceae) I. Reproduction and development of *C. racemosa* var. *laetevirens*. *Japanese Journal of Phycology/Sorui*, 35, 167–177.
- Fam`a, P., Olsen, J.L., Stam, W.T. and Procaccini, G., 2000.** High levels of intra- and inter-individual polymorphism in the rDNA ITS1 of *Caulerpa racemosa* (Chlorophyta), *European Journal of Phycology*, 35, 349–356.
DOI: 10.1080/09670260010001735951
- Fam`a, P., Wysor, B., Kooistra, W.H.C.F. and Zuccarello, G., 2002.** Molecular phylogeny of the genus *Caulerpa* (Caulerpales, Chlorophyta) inferred from the chloroplast *tufA* gene. *Journal of Phycology*, 38, 1040–1050.
DOI:10.1046/j.1529-8817.2002.t01-1-01237.x
- Fernández-García, C., Wysor, B., Riosmena-Rodríguez, R., Peña-Salamanca, E. and Verbruggen, H., 2016.** DNA-assisted identification of *Caulerpa* (Caulerpaceae, Chlorophyta) reduces species richness estimates for the eastern tropical pacific. *Phytotaxa*, 252, 185–20.
DOI: 10.11646/phytotaxa.252.3.2
- Fritsch, F.E., 1965.** The Structure and Reproduction of the Algae, Vol. I. Cambridge University Press, London. 791 P.
- Guiry, M.D. and Guiry, G.M., 2023.** AlgaeBase. World-wide electronic publication, National University of Ireland, Galway. Available at:
- John, D.M. and Al-Thani, R.F., 2015.** Benthic marine algae of the Arabian Gulf: a critical review and analysis of distribution and diversity patterns. *Nova Hedwigia*, 98(3–4), 341–392.
DOI:10.1127/0029-5035/2014/0156
- Kazi, M.A., Reddy, C.R.K. and Jha, B., 2013.** Molecular phylogeny and barcoding of *Caulerpa* (Bryopsidales) based on the *tufA*, *rbcl*, 18S rDNA and ITS rDNA genes. *PLoS ONE*, 8, e82438.
DOI:10.1371/journal.pone.0082438
- Kokabi, M. and Yousefzadi, M., 2015.** Checklist of the marine macroalgae of Iran. *Botanica Marina*, 58(4), 307–320
DOI:10.1515/bot-2015-0001

- Kumar, S., Stecher, G. and Tamura, K., 2016.** MEGA7: Molecular evolutionary genetics analysis ver. 7.0 for bigger datasets. *Molecular Biology and Evolution*, 33(7), 1870-1874
DOI:10.1093/molbev/msw054
- Leliaert, F., Verbruggen, H., Wylor, B. and De Clerck, O., 2009.** DNA taxonomy in morphologically plastic taxa: algorithmic species delimitation in the *Boodlea* complex (Chlorophyta: Siphonocladales), *Molecular Biology and Evolution*, 53, 122.
DOI:10.1016/j.ympev.2009.06.004
- Meinesz, A. and Hesse, B., 1991.** Introduction et invasion de l'algue tropicale *Caulerpa taxifolia* en Méditerranée nord-occidentale. *Oceanologia Acta*, 14, 415–26.
- Nizamuddin, M. & F. Gessner 1970.** The algae of the northern part of the Arabian Sea and of the Persian Gulf. "Meteor" Forschungsergebnisse Reihe 6: 1-42.
- Ohba, H. and Enomoto, S., 1987.** Culture studies on *Caulerpa* (Caulerpales, Chlorophyceae), II. Morphological variation of *C. racemosa* var. *laetevirens* under various culture conditions. *Japanese Society of Phycology*, 35, 178–88.
- Ohba, H., Nashima, H. and Enomoto, S., 1992.** Culture studies on *Caulerpa* (Caulerpales, Chlorophyceae) III. Reproduction, development and morphological variation of laboratory cultured *C. racemosa* var. *peltata*. *Botanical Magazine*, 105, 589–600.
DOI:10.1007/BF02489433
- Peterson, R.D., 1972.** Effects of light intensity on the morphology and productivity of *Caulerpa racemosa* (Forsskal), *Journal of Agardh Micronesica*, 8, 63–86.
- Pangestuti, R., Haq, M., Rahmadi, P. and Chun, B.S., 2021.** Nutritional Value and Biofunctionalities of Two Edible Green Seaweeds (*Ulva lactuca* and *Caulerpa racemosa*) from Indonesia by Subcritical Water Hydrolysis. *Mar. Drugs* 2021, 19(10), 578;1-17. DOI:10.3390/md19100578
- Prud'homme van Reine, W.F. and Lokhorst, G.M., 1992.** *Caulerpella* gen. nov. a nonholocarpic member of the Caulerpales (Chlorophyta), *Japanese Society of Phycology*, 40, 365–372.
DOI:10.1046/j.1529-8817.2002.t01-1-01237.x
- Riosmeña-Rodríguez, R., Ortuño-Aguirre, C. and López Vivas, J.M., 2014.** Variabilidad morfológica de *Caulerpa racemosa* (Bryopsiadales, Chlorophyta) en el Golfo de California: Implicaciones en la taxonomía. *Ciencia y Mar*, 24, 39-45.
- Salehi Balashahri1, M., Sohrabipour, J., Dehshiri, M.M. and Mehregan, I., 2021.** Molecular Identification of *Caulerpa selago* as a New Record for The Persian Gulf. *Journal of Phycological Research*, 5(2), 713-724.
DOI:10.48308/jpr.2021.102383
- Saunders, G.W., 2008.** A DNA barcode examination of the red algal family Dumontiaceae in Canadian waters reveals substantial cryptic species diversity. 1. The foliose *Dilsea*–*Neodilsea* complex and *Weeksia*. *Botany*, 86(7), 773–789.
DOI:10.1139/B08-001
- Sauvage, T., Payri, C., Draisma, S.G.A., Prud'homme van Reine, W.F., Verbruggen, H., Belton, G.S., Gurgel, C.F.D., Gabriel, D., Sherwood, A.R.**

- and Fredericq, S., 2013.** Molecular diversity of the *Caulerpa racemosa-peltata* complex (Caulerpales) in New Caledonia, with new Australasian records for the variety cylindracea. *Phycologia*, 52, 6–13.
DOI:10.2216/11–116.1
- Senerpont Domis, L.N.D.E., Fama', P., Bartlett, A., Prud'homme van Reine, W.F., Armenta Espinosa, C. and Trono G.C.J.R., 2003.** Defining taxon boundaries in members of the morphologically and genetically plastic genus *Caulerpa* (Caulerpales, Chlorophyta), *Journal of Phycology*, 39, 1019–1037.
DOI:10.1111/j.0022-3646.2003.02-203.x
- Shams, M. and Amini, N.G., 2017.** A New Taxonomic Survey of *Caulerpa* Lamouroux Species (Chlorophyta, Caulerpales) in the Southern Coasts of Iran. *American Academic Scientific Research Journal for Engineering, Technology, and Sciences*, 36(1), 45–57.
- Sohrabipour, J. and Rabiei, R., 2007.** The checklist of green algae of the Iranian coastal lines of the Persian Gulf and Gulf of Oman. *Iranian Journal of Botany*, 13, 146–149.
DOI:20.1001.1.1029788.1386.13.2.17.7
- Sohrabipour, J. and Rabiei, R., 1996.** New records of algae from Persian Gulf and flora of Iran. *Iranian Journal of Botany*, 7, 95–115.
DOI:10.22092/IJB.2015.103483
- Sohrabipour, J. and Rabiei, R., 1999.** A list of the algae of sea shores of Persian Gulf and Oman Sea in the Hormozgan Province. *Iranian Journal of Botany*, 8, 131–162.
DOI: 20.1001.1.1029788.1378.8.1.16.7
- Stam, W.T., Olsen, J.L., Zaleski, S.F., Murray, S.N., Brown, K.R., 2006.** A forensic and phylogenetic survey of *Caulerpa* species (Caulerpales, Chlorophyta) from the Florida coast, local aquarium shops, and ecommerce: establishing a proactive baseline for early detection. *Journal of Phycology*, 42, 1113–1124.
DOI:10.1111/j.15298817.2006.00271.x
- Verlaque, M., Boudouresque, C.F., Meinesz, A. and Gravez, V., 2000.** The *Caulerpa racemosa* complex (Caulerpales, Ulvophyceae) in the Mediterranean Sea. *Botanica Marina*, 43, 49–68.
DOI:10.1515/BOT.2000.005
- Verlaque, M., Durand, C. H., Huisman, J.M., Boudouresque, Ch. F. and Le Parco Y., 2003.** On the identity and origin of the Mediterranean invasive *Caulerpa racemose* (Caulerpales, Chlorophyta), *European Journal of Phycology*, 38, 325–339.
DOI:10.1080/09670260310001612592
- Wirawan, I.G.P., Dewi, N.K.E.S., Sasadara, M.M.V., Sunyamurthi, I.G.N.A., Jawi, I.M., Wijaya, N., Darmawati, I.A.P., Suada, I.K. and Krisnandika, A.A.K., 2022.** Phytochemical Analysis and Molecular Identification of Green Macroalgae *Caulerpa* spp. from Bali, Indonesia. *Molecules*, 27, 4879.
DOI:10.3390/molecules27154879