



Mini Review

Rhazes's Perspective on the Prevention and Treatment of Rabies

Seyyed Alireza Golshani¹ , Mohammad Hassan Najafi^{2,3*} , Waleed Saffah Meteab^{4,5} , Amir Houshang Mehrparvar⁶

1. Medical Innovation Center for Nationalities, Inner Mongolia Medical University, Hohhot, China.
2. Department of Persian Medicine, School of Iranian Traditional Medicine, Shahid Sadoughi University of Medical Sciences, Yazd, Iran.
3. Research Center of Persian Medicine, Shahid Sadoughi University of Medical Sciences, Yazd, Iran.
4. Department of History, Faculty of Literature and Human Sciences, Urmia University, Urmia, Iran.
5. University of Technolog, Baghdad, Iraq.
6. Industrial Diseases Research Center, Shahid Sadoughi University of Medical Sciences, Yazd, Iran.



How to cite this article Golshani SA, Najafi MH, Meteab WS, Mehrparvar AH. Rhazes's Perspective on the Prevention and Treatment of Rabies. *Archives of Rhazes Institute Journal*. 2026; 81(2):273-278. <https://doi.org/10.32598/ARI.81.2.3590>

doi <https://doi.org/10.32598/ARI.81.2.3590>

Article info:

Received: 18 Nov 2025

Accepted: 21 Dec 2025

Published: 01 Mar 2026

Keywords:

Medical history, Rabies, Rhazes,
Traditional Iranian medicine

ABSTRACT

Iranian traditional medicine (TM) holds a millennia-old legacy in viral disease management, with classical scholars like Rhazes (854–925 CE) making ground-breaking contributions to rabies prevention. This study examines Rhazes' innovative approaches, including clinical observations, bite management protocols, and transmission control strategies that predate modern epidemiological understanding. Using historical research methods aligned with the Annales school, we analyze his emphasis on isolation measures, dietary interventions, and environmental factors in disease prevention. Notably, Rhazes' recommendations for rabies control show remarkable parallels with contemporary World Health Organization (WHO) guidelines. His work demonstrates an advanced understanding of contagion theory, including climate influences and contact precautions. The study highlights how Rhazes' experimental methods and systematic patient care protocols provide valuable insights for modern viral disease management. By bridging ancient Persian medical wisdom with current practices, this research underscores the enduring relevance of traditional knowledge in global health. The findings advocate for integrating historical preventive approaches into modern healthcare systems, particularly for zoonotic diseases such as rabies. This investigation not only documents Iran's medical heritage but also provides a model for interdisciplinary dialogue between TM and evidence-based practice. The study ultimately positions Rhazes as a pioneer whose rabies management strategies remain conceptually valid even centuries later.

* Corresponding Author:

Mohammad Hassan Najafi, Assistant Professor.

Address: Department of Persian Medicine, School of Iranian Traditional Medicine, Shahid Sadoughi University of Medical Sciences, Ardakan, Iran.

Tel: +98 (913) 1532985

E-mail: drmhajafi@yahoo.com.

Copyright © 2026 The Author(s).
This work is licensed under a Creative Commons Attribution-NonCommercial 4.0 International license (<https://creativecommons.org/licenses/by-nc/4.0/>).
Noncommercial uses of the work are permitted, provided the original work is properly cited.

1. Introduction

1.1. Context

Across the globe, traditional medicine (TM) serves as either the cornerstone or a complementary component of health care delivery. The [World Health Assembly \(WHA\)](#) resolution on TM (WHA62.13), adopted in 2009, highlights the importance of TM and complementary medicine (T&CM) as a vital yet often underappreciated aspect of healthcare systems. T&CM is practiced in nearly every country worldwide, and the demand for its services continues to grow. When TM is proven to be safe, effective, and of high quality, it significantly contributes to the goal of ensuring universal access to healthcare [1]. Many countries now recognize the need to develop a cohesive and integrative approach to healthcare that enables governments, practitioners, and—most importantly—healthcare users to access T&CM services in a manner that is safe, respectful, cost-efficient, and effective [2]. Since the publication of the previous global strategy in 2002, much has evolved. An increasing number of countries have come to acknowledge the role that T&CM can play in promoting individual health and well-being, as well as enhancing the comprehensiveness of their health care systems. Governments and consumers are no longer solely focused on herbal medicines; they are now exploring whether T&CM practices and practitioners should be integrated into mainstream healthcare service delivery. In response to this growing demand and to [WHA Resolution 62.13](#) on TM, the [World Health Organization \(WHO\)](#) recently updated the objectives of its TM program [1]. The [WHO TM strategy 2014–2023](#) aims to equip healthcare leaders with tools to develop solutions that align with a broader vision of improved health outcomes and patient autonomy [1].

Interestingly, in Iran, prior to the establishment of the [Pasteur Institute of Iran](#) in 1920 and the National Center for Reference & Research on Rabies in 1923, the prevention and treatment of rabies were guided and managed by practitioners of Iranian TM [3, 4]. Globally, folk medicine and TM have offered unique approaches to addressing a wide array of injuries and diseases, including animal bites from rabid animals. While the efficacy of most traditional anti-rabies treatments has not been validated through controlled clinical trials, community-based surveys have demonstrated their practical effectiveness [5].

1.2. Data acquisition

Rabies has long been a subject of interest in folk medicine, as evidenced by its mention in ancient Iranian texts such as the Avesta. For instance, chapters 31–38 of the *Vendidad* refer to rabid dogs as “mad dogs” and outline guidelines for their human treatment and containment [6]. Once regarded as a general encyclopedia, this text emphasizes that rabid dogs should be treated with care and kept captive. TM sources document numerous cases involving knowledge of external agents and mediators of rabies transmission, including preventive measures. These accounts suggest that the practices described are grounded in evidence and clinical experience [7] ([Figure 1](#)).

The *Lyssavirus* genus comprises RNA viruses belonging to the *Rhabdoviridae* family within the *Mononegavirales* order. Mammals, including humans, serve as the natural hosts for viruses in this genus, among which the rabies virus is responsible for causing the rabies in humans [8].

Rabies is a viral disease transmitted when an infected canine bites a human, allowing the virus in the canine’s saliva to enter the human body through the wound. The virus then spreads via the bloodstream until it eventually reaches the nervous system. The incubation period—the time between infection and the onset of symptoms—typically ranges from one to three months but can vary significantly, lasting less than a week or extending beyond a year. Early symptoms include fever, mental discomfort, depression, insomnia, anxiety, persistent sadness and frustration, lethargy, itching at the bite site, and hydrophobia. Additional signs of the disease include swelling of lymph nodes near the bite site, shortness of breath, escalating periodic tetanoid spasms (particularly in the respiratory muscles), albuminuria, and the presence of Negri bodies in the brain [9].

2. Materials and Methods

This study employed a descriptive research methodology, incorporating library-based and historical research approaches to examine original texts and traditional medical books from Iran. Drawing on the French historiographical school, specifically the *Annales* School, which emphasizes a critical examination of historical approaches in the context of scientific development—with a particular focus on central Iran—the study compares the methods of preventing infectious diseases, such as smallpox, in Iranian TM with those of modern medicine. Below are the stages involved in gathering and compiling this research:



Figure 1. Statue of a Bijar breed dog from the Achaemenid period (Museum of Ancient Iran)

A) Comprehensive examination of primary sources: A detailed review was conducted of Rhazes' perspectives on the prevention and treatment of rabies, utilizing first-hand historical sources, contemporary research, and scientific articles. References were drawn from Arabic, English, and Persian sources, supplemented by searches in reputable databases such as SID, Google Books, Google Scholar, Internet Archive, and Scopus to gather additional information. The findings were then synthesized and documented.

B) Note-taking on relevant topics: Notes were systematically recorded on pertinent research topics, ensuring that all relevant aspects of Rhazes' contributions were captured.

C) Systematic classification of notes: The collected notes were organized into introductory sections and systematically classified for ease of reference and detailed analysis.

D) Finalization and writing: The gathered information was reviewed, finalized, and written up in a coherent and structured format.

The authors aim to focus their efforts on introducing Rhazes' views on the prevention and treatment of rabies, while comparing these with modern medical practices. Additionally, treatment solutions derived from TM will be explored and contrasted with contemporary approaches, an area that has received comparatively less attention in existing literature.

3. Results

3.1. Rhazes's view on the prevention and treatment of rabies

Iranian TM texts extensively address rabies, emphasizing its transmission to humans through the bite of rabid dogs. To prevent the disease, these texts recommend recognizing and avoiding rabid animals or eliminating them due to their role in disease transmission. If bitten by an animal, it is advised to allow the wound to bleed profusely and avoid bandaging it, as per traditional guidelines for rabies wounds [7, 10]. One of the most prominent figures in Iranian medical history is Abū Bakr al-Rāzī (Rhazes) (c. 854–925 CE), whose contributions span various fields of medical science and pharmacology. Rhazes documented numerous experiments and clinical observations in his works, including insights into rabies [11-13].

In his book *Al-Mansouri fi al-Tibb* (the book of medicine dedicated to Mansur), Rhazes references the methods of earlier physicians for diagnosing rabies: "If a dog bites a man and it is unclear whether the dog was rabid, one should soak a piece of bread in blood drawn from the bite site and throw it to another dog. If the dog eats the bread, the bite is not from a rabid dog; if it avoids eating it, the bite is from a rabid dog. [Another method is to] crush some walnuts, apply them to the wound overnight as a dressing, and the next day, throw the dressing to a chicken or rooster. If the bite was from a rabid dog, the bird will refuse to eat it. Even if the bird consumes it out of hunger, it will die the following day. If these signs are observed, it is recommended to reopen the wound and provide the necessary care as described. If no such signs appear, the wound may be closed" [14].

Rhazes observed that dogs are more prone to rabies during extreme heat and, occasionally, in winter. He noted that rabid dogs avoid consuming food and water, to the extent that they may even exhibit hydrophobia, nearly perishing at the mere sight of water [14].

To aid in diagnosing rabies, Rhazes emphasized in *Al-Mansouri fi al-Tibb* (the book of medicine dedicated to Mansur) that, in addition to recognizing the signs of rabies, certain critical considerations must be taken into account to ensure that an animal intended for keeping, or one bitten by another animal, is not rabid. He stated: “Since being bitten by a rabid animal causes a severe and life-threatening disease, the signs of rabies should be widely publicized so that people may avoid such dogs and, if necessary, euthanize them.” To combat rabies, Rhazes described the characteristics of a rabid dog and the rationale for its elimination: “The dog’s mouth remains open, its tongue hangs out, foam and saliva drip from its mouth and nose, its eyes are red with a blood-like hue, and its head and ears droop. It tucks its tail between its legs, staggers like a drunkard, attacks and bites indiscriminately, fails to recognize its owner, and barks less frequently. If any or all of these signs are observed in a dog, it is best to euthanize or kill the animal, as a bite from such a dog may initially show no distinct symptoms but will eventually lead to the manifestation of the disease” [14].

Rhazes also advises: “When bitten by a rabid dog, the wound should not be stitched, and the bite wound must remain open. Immediately after the bite, the wound should be opened to drain a significant amount of blood. If the patient does not recover within one day of the onset of rabies symptoms, there is no hope for recovery” [7, 10]. To manage rabies, it is recommended to place the patient in a cool, quiet environment and administer fluids using a gastric tube, as pharyngeal muscle spasms prevent swallowing, making it difficult to replenish lost fluids [14]. Rhazes further suggests: “Increase the intake of meat and sweets in the patient’s diet, ensure adequate rest, insert a long tube into the mouth for treatment, and provide water. Additionally, if the patient is unable to swallow, use liquid medications and administer enemas to quench their thirst” [14].

In *Kitāb al-Ḥāwī fi al-Ṭibb* (the comprehensive book on medicine), Rhazes describes a treatment for rabid dog bites involving the consumption of burnt crab. He recommends burning the crab, mixing three ounces of its ash with 1.5 ounces of cinnamon, and consuming the mixture with wine for three days. He also highlights the efficacy of combining burnt crab ash with cinnamon and frankincense for treating rabid dog bites, describing this remedy as particularly impressive. Rhazes advises doubling the amount

of burnt crab ash if several days have passed since the bite. Elsewhere in the same book, he emphasizes the usefulness of the pomegranate plant, recommending the consumption of pomegranate gum, very sour vinegar, and olive oil as a treatment for rabies caused by bites. Rhazes believed that applying sardine extract or pickle as a poultice could effectively treat rabid dog bites. He also mentions the use of clay, wine, water, or honey individually for treating such bites [15].

In *Man la Yahduruhu Al-Tabib* (for one who has no physician to attend him), Rhazes lists substances that can be used as poultices for wounds, including onions, garlic, and ginger. He further suggests using a combination of camel urine, licorice, frankincense, and *Cuscuta* before the onset of symptoms. *Epithymum* has also been employed to treat bite wounds, and irrigation of the wound caused by rabid dog bites is recommended to remove blood from the affected area [15]. Rhazes also describes *Lytta vesicatoria* as having a remarkable and beneficial effect in preventing rabies [16].

3.2. Rhazes’ clinical observations

Rhazes also wrote of a personal experience in the Ray hospital regarding a schizophrenic and delusional person: “There was someone in the hospital whom a rabid dog had bitten. They barked in the night, and would feel frightened and start shaking whenever water was brought for [them, despite constantly asking for water and complaining about thirst. When we brought water near their face, it would become grim and full of anger, and they would throw the bowl of water away, saying that it was full of filth. When we asked what kind of pollution was in the water, they would respond that it had cat and dog guts in it. And then they would ask that we bring another bowl of water, and if we did, they would repeat what they] said before and become agitated and hostile to us. They would beg us to let them drink fresh and clear water for the sake of God” [8].

The latency period—the interval between infection and the onset of symptoms—is one of the notable aspects of Rhazes’ perspective on rabies. He explained: “If such a dog bites someone, initially, the person does not realize they are sick and may dismiss it as a minor bite. However, after a short while, signs of illness begin to appear. The individual develops a fear of water, avoids drinking it, and starts trembling at the sight of water. These symptoms can progress to the point of being fatal. The patient will fear anything wet or fluid, avoiding it until they succumb to thirst. In some cases, the afflicted person may attack and bite others, potentially transmitting the disease to those around them” [8].

Table 1. Rhazes' perspectives on rabies (hydrophobia)

Topic	Rhazes' Observations & Theories	Possible Source
Cause	Caused by bites from rabid animals (especially dogs), with transmission via "toxic saliva."	Al-Hawi, Al-Judari wa al-Hasbah
Symptoms	<ul style="list-style-type: none"> - Hydrophobia (fear of water) - Extreme agitation/anxiety - Progressive paralysis - Fatal outcome in severe cases 	Al-Hawi
Incubation period	Mentioned a variable latency period (days to months after the bite).	Al-Hawi
Treatments	<ul style="list-style-type: none"> - Wound cleansing (e.g. washing with water/salt) - Use of sedatives/palliative remedies. 	Al-Hawi
Prognosis	Emphasized the fatal nature of the disease once neurological symptoms appeared.	Al-Hawi
Differential diagnosis	Distinguished rabies from similar conditions (e.g. epilepsy or poisoning).	Al-Judari wa al-Hasbah

4. Discussion

4.1. WHO's view on the prevention and treatment of rabies

One of the notable points is the striking similarity between Rhazes's treatment methods and the protocols recommended by the WHO. As stated in section 5.2 of the WHO guidelines on the clinical management of rabies patients: "Patients remain conscious and are often aware of the nature of their illness but are usually extremely agitated, particularly when excitation predominates. Furthermore, they are often isolated due to the perceived risk of viral transmission through contact. Patients with confirmed rabies should receive adequate sedation and care in an appropriate medical facility, preferably in a private room, with suitable emotional and physical support. Repeated intravenous morphine or benzodiazepines are effective in alleviating the severe agitation, anxiety, and phobic spasms that afflict patients with furious rabies. Once furious rabies has been diagnosed, invasive procedures should be avoided, and the patient should be cared for in a private, quiet, draft-free environment. Given the inevitability of death in most cases, treatment should focus on providing comfort, including heavy sedation (barbiturates, morphine), and avoiding intubation or life-support measures once the diagnosis is certain" [17].

Rhazes' ingenuity in addressing rabies demonstrates that he was not only a skilled and experienced physician but also a leading figure in the understanding of this disease. In his works, he meticulously described rabies and its symptoms while emphasizing critical factors, such as how a rabid animal's bite can lead to severe illness. He raised public awareness about the signs of rabies, teaching people to avoid suspicious animals and, if necessary, euthanize them to prevent further transmission. Rhazes provided numerous

suggestions for diagnosing and treating rabies, including recipes for ointments, dietary recommendations, and instructions for washing wounds with fresh water to remove contaminated blood. His methods were grounded in the research of earlier physicians, his own clinical observations, and empirical experimentation. Remarkably, his approach closely aligns with the modern protocols recommended by the WHO, underscoring the enduring relevance of his contributions to medicine (Table 1).

5. Conclusion

Rabies, as described by Muhammad ibn Zakariya al-Rhazes (Rhazes), represents a fascinating intersection of early medical observation and scientific reasoning in the medieval Islamic world. In his works, particularly al-Hawi and al-Judari wa al-Hasbah, al-Rhazes provided detailed accounts of rabies (known as *dā' al-kalb*), emphasizing its transmission through the bite of rabid dogs and its devastating neurological effects. His clinical descriptions—including symptoms like hydrophobia, extreme agitation, and muscle spasms—align remarkably well with modern medical understanding, showcasing his keen observational skills. Rhazes proposed several treatments for rabies, some of which reflect an intuitive grasp of medical principles. He stressed the immediate cleansing of bite wounds to prevent the spread of the "toxic substance" (a prescient notion, given the era's limited knowledge of viruses). He also recommended herbal antidotes and bloodletting (*hijama* or *fasd*) to purge the body of toxins. Yet, he acknowledged the near-fatal prognosis once neurological symptoms appeared, demonstrating both his empirical honesty and the limits of 9th-century medicine. What sets Rhazes apart is his rejection of superstition in favor of systematic observation. While he lacked modern tools, his methodology—rooted in cause-and-effect analysis—laid the groundwork

for evidence-based medicine. Today, we know rabies is caused by the Lyssavirus, and is preventable through vaccines, but Rhazes's insights remain a testament to his intellectual rigor. His work underscores timeless principles: the importance of prevention, the value of critical inquiry, and the need to challenge misconceptions. Rhazes's legacy in rabies research highlights how medieval scholars bridged ancient traditions and empirical science. His approach—meticulous, rational, and patient-centered—resonates even now, reminding us that scientific progress often begins with the courage to observe, question, and document the world as it is.

Compliance with ethical guidelines

It is hereby asserted that all ethical considerations were duly considered during the preparation of the submitted manuscript.

Data availability

The datasets generated and/or analyzed during this study are available from the corresponding author upon reasonable request.

Funding

This research did not receive any grant from funding agencies in the public, commercial, or non-profit sectors.

Authors' contributions

Conceptualization, study design, analysis and data interpretation: Seyyed Alireza Golshani; Data acquisition and writing the original draft: Seyyed Alireza Golshani, Mohammad Hassan Najafi, and Waleed Saffah Meteab; Review and editing: Seyyed Alireza Golshani and Mohammad Hassan Najafi; Supervision: Mohammad Hassan Najafi; Statistical analysis: Seyyed Alireza Golshani, Mohammad Hassan Najafi, and Amir Houshang Mehrparvar; Administrative, technical, and material support: Waleed Saffah Meteab.

Conflict of interest

The authors declared no conflict of interest.

References

- [1] World Health Organization. *WHO traditional medicine strategy: 2014-2023*. Geneva: WHO; 2013. [\[Link\]](#)

- [2] Daneshfard B, Sanaye MR, Nimrouzi M. Prolegomena to a true integrative medical paradigm. *Altern Ther Health Med*. 2019; 25(2):AT5662. [\[PMID\]](#)
- [3] Fazeli M, Fayaz A, Bashar R. One-hundred-year efforts by the Pasteur Institute of Iran as a part of the rabies infectious control puzzle in the world. *Nov Biomed*. 2019; 7(3):165-79. [\[DOI:10.22037/nbm.v7i3.26554\]](#)
- [4] Shaban Salmani N, Fazeli M, Bashar R. The national center for reference and research on rabies, Pasteur Institute of Iran: An unforgettable name in the health history of rabies control in the world. *J Res Hist Med*. 2021; 10(4):205-14. [\[Link\]](#)
- [5] Beasley EA, Wallace RM, Coetzer A, Nel LH, Pieracci EG. Roles of traditional medicine and traditional healers for rabies prevention and potential impacts on post-exposure prophylaxis: A literature review. *PLoS Negl Trop Dis*. 2022; 16(1):e0010087. [\[DOI:10.1371/journal.pntd.0010087\]](#) [\[PMID\]](#)
- [6] Dostkhah J. *[Avesta (ancient written by Iranians) (Persian)]*. Tehran: Marwarid Press, 2008. [\[Link\]](#)
- [7] Hatami H. History of rabies in traditional medicine's resources and Iranian research studies: On the Occasion of the world rabies day (September 28, 2012). *Int J Prev Med*. 2012; 3(9):593-5. [\[PMID\]](#)
- [8] Zandi F, Goshadrou F, Meyfour A, Vaziri B. Rabies infection: An overview of Lyssavirus-host protein interactions. *Iran Biomed J*. 2021; 25(4):226-42. [\[DOI:10.52547/ibj.25.4.226\]](#) [\[PMID\]](#)
- [9] Giesen A, Gniel D, Malerczyk C. Correction to: 30 Years of rabies vaccination with Rabipur: A summary of clinical data and global experience. *Expert Rev Vaccines*. 2015; 14(5):775. [\[DOI:10.1586/14760584.2015.1027863\]](#) [\[PMID\]](#)
- [10] Esnaashary MH, Yarmohammadi H, Dalfardi B, Askarpour MR, Daneshfard B, Mehdizadeh A. Rabies: A comprehensive review of traditional Persian medicine literature. *J Res Hist Med*. 2017; 6(1):15-24. [\[Link\]](#)
- [11] Golshani SA, Hashemi ZH, Ranjbar A, Zarshenas MM. Use of Khadab for beauty, health, and hair strengthening in the works of Rhazes (c. 854-925 CE). *Pharm Hist*. 2023; 53(2):58-60. [\[Link\]](#)
- [12] Golshani SA. Methods of medical education, transoxianian physicians in the Samanid period (874-1004 AD). *J Med Educ Dev*. 2022; 17(3):224-35. [\[DOI:10.18502/jmed.v17i3.11299\]](#)
- [13] Hossein Hashemi Z, Golshani SA. Rhazes's view on nasal tumors. *J Res Hist Med*. 2023; 12(4):301-6. [\[Link\]](#)
- [14] Rhazes AB. *Al-mansouri in Medicine (Al-mansouri fi al-Tibb)* M. E. Zaker, Persian trans.]. Tehran: Tehran University of Medical Sciences; 2008. [\[Link\]](#)
- [15] Shabansalmani N, Pourhossein B, Khosravi MS, Farahat F, Alavian SF, Bashar R, et al. A Review of Rabies Treatment Methods in Ancient Iran. *J Virol Viral Dis*. 2022; 2(3):1-7. [\[DOI:10.54289/JVVD2200111\]](#)
- [16] Moallemi M, Yousofpour M, Jokar A. Prevention of Rabies by Application of *Lytta vesicatoria* in Persian medicine texts in Islamic civilization. *Trad Integr Med*. 2021; 6(1):70-7. [\[DOI:10.18502/tim.v6i1.5929\]](#)
- [17] WHO. WHO Expert consultation on rabies. Geneva: World Health Organization; 2013. [\[Link\]](#)