



Total phenolic and flavonoid content of *Satureja sahandica* plant under the influence of animal manure and enriched straw in dry conditions

Seyedhadi Hosseini^a, Khalil Karimzadeh Asl^b, Hamid Mozafari^a, Behzad Sani^a and Mahdi Mirza^b

^a Department of Agronomy, Shahr-e-Qods Branch, Islamic Azad University, Tehran, Iran.

^b Faculty of Research Institute of Forests and Rangelands, Agricultural Research, Education and Extension Organization (AREEO), Tehran, Iran.

ARTICLE INFO	ABSTRACT
Corresponding Author: Seyedhadi Hosseini Seyedalhosseini63@yahoo.com Received: 22 April 2024 Accepted: 13 May 2024 Keywords: Antioxidant enzymes Dry farming conditions Flavonoid Phenol <i>Satureja sahandica</i>	<i>Satureja sahandica</i> is an exclusive species of Iran. The aerial parts of this plant are widely used in food and pharmaceutical industries. Soil fertility management is one of the main factors determining the growth and quantitative and qualitative performance of medicinal plants. According to the statistics presented in 2024, Iran is in the year of drought that it multiplies the necessity of using the climate capacity correctly. Agriculture in dry farming conditions is very important considering the available water capacities in the world. Because something like 41% of the world includes dry land. These experiments were carried out in the form of as split plots in the form of a randomized complete block design in the steppe region (Humand area of Damavand city). The experimental treatments included organic manure (cow, straw enriched with ammonium sulfate and control treatment), planting density (low density of 26666 plants per hectare, medium density of 40000 plants per hectare and high density of 80000 plants per hectare) in 2017 and 2018. Sahandi Savory seeds were obtained from the Research Institute of Forests and Rangelands of Iran (RIFR). They were planted in culture trays filled with a mixture of peat moss and vermiculite (1:1 volume) under greenhouse conditions. Two-month-old seedlings were brought to the farm from the greenhouse in April 2017. The average annual rainfall was 333 mm. This research showed that increasing the soil fertility can make a significant change in the increase of phenol and flavonoids. Copyright © 2022 Union Medicinal Plants of Iran. All rights reserved.

1. Introduction

Savory (*Satureja sahandica*) is an exclusive species of Iran (Yousefzadi et al., 2012). The aerial parts of this plant are widely used in food and pharmaceutical industries and can replace artificial preservatives as a natural antioxidant (Hajhashemi et al., 2002). The vegetative growth of this Sahandi spice starts from the end of March and its flowering continues from the middle of June to the end of August. If it is harvested or trapped, its flowering continues until early autumn (Akbarinia et al., 2009). Harvesting medicinal plants at the wrong time not only reduces the amount of the obtained product, but also the harvested product will not have the desired quality. Because the function of the target organ as well as the amount of secondary metabolites of a medicinal plant is different in different stages of plant growth and development (Omidbaigi, 2005). Plants have a set of antioxidants to control the amount of free oxygen radicals

and to protect their plant cells in stressful conditions (Borsani et al., 2001). Biological activities of plant extracts, especially their anti-aggregation and free radical activities (Brunetti et al., 2013) as well as antimicrobial activity and capacities (Pistelli & Giorgi., 2012) and anticancer (Batra & Shama., 2013) Due to the presence of complex compounds, they contain phenol and flavonoids (Gopčević et al., 2019). Phenols are a large group of biologically active compounds (over 8000 compounds) that are among the ubiquitous secondary metabolites in plants (Dreosti, 2000). Phenols are classified into two basic groups, simple phenols and polyphenols, based on the number of phenol subunits (Marinova et al., 2005). The group of simple phenols contains phenolic acids or phenols with a carboxyl group, which determines their functional properties. There are at least two phenol rings in polyphenol. Flavonoids belong to this group. More than 4000 flavonoids have been identified in different



plant species (Harborne & Tumer., 1984). The integrated management of plant nutrition is called an operation that uses nutrients, water, products and plant cover management methods in the soil bed, according to the cultivation and the specific agricultural system, with the aim of improving and preserving soil fertility, land productivity and reducing destruction in environment is done (Mohebi, 2016). The integrated management of nutrients is done with the aim of optimizing soil conditions, according to its chemical, physical, biological and hydrological properties, in order to increase the productivity of the farm, so that soil degradation is as low as possible (Feller et al., 2012). Soil fertility management is one of the main factors determining the growth and quantitative and qualitative performance of medicinal plants (Omidbaygi., 1995). In high densities, the competition for moisture, food and light increases, which results include the reduction of stem diameter and the increase of plant height (Gozubenli et al., 2003). One of the important factors to obtain the highest seed yield in corn is to determine the appropriate density according to the climatic conditions of each region and the characteristics of the cultivars to be planted (Akintoye et al., 1997). The changing pattern of the hydrological cycle is one of the issues related to climate change that deserves urgent attention (Donat et al., 2016). Evaporation, transpiration is related to water and soil balance and has a key role in the interactions of climate, soil and vegetation (Yang et al., 2007). Water management in dry farming agriculture systems is important in addition to providing the water needed for food production, in order to create resistance to the risks and uncertainties related to water use. The availability of water is different in different regions, and the competition to obtain water can become very intense in regions where natural rainfall decreases, such as arid and semi-arid regions (Lovelli, 2019). Rainfed agriculture and dry farming agriculture are terms that are often used as synonyms, but this is a mistake, in both cases irrigation is omitted, but by omitting this aspect, their meaning is profoundly different (Stewart., 2016,a). But the latest methods are to conserve water and the sustainable performance of crops, limit fertilizers and other inputs and act to limit erosion (Stewart, 2016, b). Agriculture in dry farming conditions is very important considering the available water capacities in the world, because something like 41% of the world includes dry lands (Lovelli, 2019) that In order to develop and cultivate them, we should turn to dry farming agriculture.

2. Materials and Methods

This experiment was carried out based on split plots in the form of randomized complete blocks design in the Humand area of Damavand city. The experimental

treatments included organic manure (cow, straw enriched with ammonium sulfate and control treatment), planting density (low density of 26666 plants per hectare, medium density of 40000 plants per hectare and high density of 80000 plants per hectare) in 2017 and 2018. Sahandi Savory seeds were obtained from the Research Institute of Forests and Rangelands of Iran (RIFR). They were planted in culture trays filled with a mixture of peat moss and vermiculite (1:1 volume) under greenhouse conditions. Two-month-old seedlings were brought to the farm from the greenhouse in April 2017. The average annual temperature of the farm is 11°C, minimum -24°C in January-February and maximum 37°C in July-August. The average annual rainfall was 333 mm. Soil pH and EC (0-20 cm) were 8.3 and 0.8 dsm⁻¹, respectively (Table 1). The results of the physicochemical analysis of the soil of savory planting bed in dry conditions are as described in Table 1.

Table.1-Physico-chemical analysis of the soil of savory planting bed in dry conditions


Characteristics	Values
clay (percent)	5/35
sand (percent)	17
silt (percent)	5/47
soil pattern	Rosy Silty
CEC (m.e/100 gr)	26
soil acidity	48/7
Electrical conductivity (decsiemens/meter)	0.522
organic carbon (percentage)	0.98
Total nitrogen (percentage)	0.072
Absorbable potassium (mg/kg)	800

Table.2- Meteorological information of planting area

Year	Average temperature (° C)	Annual rainfall (mm)
2016	12.38	321.5
2017	11.95	487

In order to improve soil fertility, the consumption of rotted cow manure required for the implementation of the project was determined based on 30 tons per hectare. To apply fertilizer treatment, we took 36 kg of rotted cow manure for each sub-plot and spread it evenly in the created furrows. Then we covered the manure with pile soil. The amount of straw needed to implement each plan was based on 10 tons per hectare. In order to carry out the treatment of enriched straw, the consumption of 12 kilograms of chopped wheat straw was uniformly distributed in the created grooves of each sub-plot. And the consumption of 240 grams of ammonium sulfate (based on 2 kilograms for every 100 kilograms of straw in the silo) after dissolving in 20 liters of water was sprayed uniformly on the straws of the floor. Then it was covered with pile soil. After applying the desired treatments and planting, annual statistics were taken. Phenolic compounds are measured by methods that use Folin-Ciocalteu as a reagent and gallic acid as a standard. Phenolic content was measured by spectrophotometer,

according to Ouchikh (2011) method. In this experiment, 2 grams of the sample was homogenized with 8 milliliters of 80% ethanol and centrifuged at 12,000 x g for 20 minutes, and then 0.5 milliliters of the supernatant was removed using a sampler and poured into 15 milliliter flasks. , then 500 microliter Folin-Ciocalto was added to the contents of Falcon and after 2 minutes, one milliliter of 7% sodium carbonate was added to the reaction mixture and the final volume was brought to 6 milliliters using distilled water. The falcons were placed in a 30°C bain-marie bath (dark conditions) for 90 minutes. The absorption of the samples was also measured in a spectrophotometer at a wavelength of 725 nm. This method was used for all gallic acid standard solutions and standard calibration curve drawing. The amount of flavonoid was measured by aluminum chloride colorimetric method in this method, 0.5 ml of extract solution with 1.5 ml of 95% ethanol, 0.1 ml of 10% aluminum chloride, 0.1 ml of 1 M potassium acetate and 2.8 ml a liter of distilled water was mixed. After keeping the samples at room temperature for 30 minutes, the absorbance of the mixture was read at a wavelength of 415 nm. Querstein standard was used to draw the curve (Chang, et al., 2002). The amount of enzyme activity was measured by the method of Ranieri et al. (2003). As a result of the reaction between ascorbate peroxidase and ascorbic acid and H₂O₂, dehydroascorbate is produced, which is read at a wavelength of 290 nm. The reaction medium contained 600 microliters of 0.1 mm EDTA, 1500 microliters of 50 mm phosphate buffer (pH = 7), 400 microliters of 0.5 mm ascorbic acid, 400 microliters of 30% H₂O₂, and 50 microliters of enzyme extract. Enzyme activity measurement was recorded for 7 minutes with 20 second intervals. Ascorbate peroxidase also converts H₂O₂ into water using ascorbate, which acts as an electron donor, according to the following equation:




The specific activity of the enzyme was reported as micromoles of oxidized ascorbate per minute per milligram of protein.

$$\text{Activity (U/ml)} = \frac{\Delta A_{290} \times l \times V_t \times d_f}{\varepsilon \times l \times t \times V_s}$$

μmol ascorbate oxidized min⁻¹ mg⁻¹ protein.

The activity of catalase enzyme (CAT) was measured at 25 degrees Celsius using a spectrophotometer and according to the method of Aebi (1984). A spectrophotometer made in Japan at a wavelength of 240 nm was used to measure enzyme activity. The solutions and materials used included 3000 microliters of phosphate buffer (pH = 7) 50 millimolar, 5 microliters of 30% H₂O₂ and 50 microliters of enzyme extract. And enzyme activity was recorded for 5 minutes in 20 second intervals. Catalase enzyme converts H₂O₂ into O₂ and

H₂O according to the equation below without the need for a regenerating agent.



The specific activity of the enzyme was reported as micromoles of H₂O₂ dissolved per minute per milligram of protein.

$$\text{Activity (U/ml)} = \frac{\Delta A_{290} \times l \times V_t \times d_f}{\varepsilon \times l \times t \times V_s}$$

μmol of H₂O₂ decomposed min⁻¹ mg⁻¹ protein.

Superoxide dismutase activity was measured according to the method of Dhindsa et al. (1981). The activity of this enzyme is evaluated photo metrically. The main reaction buffer included 100 mm phosphate buffer (pH=7.8), 12 mm methionine, 75 μm nitro blue tetrazolium, 100 μm EDTA, and 0.025% Triton X-100. 290 microliters of the main buffer was added to each well. Then 5 μM of 2 μM riboflavin buffer was added to the reaction mixture and the device was calibrated at a wavelength of 560 nm. 10 microliters of protein extract was used to measure each sample. This reaction is evaluated based on the rate of photo regeneration of nitro blue tetrazolium and the ability of superoxide dismutase enzyme to inhibit this reaction. In order to measure polyphenol oxidase, 0.05 g of fresh leaf tissue was homogenized in a cold Chinese mortar and in an ice container with 2 ml of 0.1 M phosphate buffer with an acidity of 6.8 and then it was centrifuged for 15 minutes at 13,000 revolutions per minute at a temperature of 4 degrees Celsius. The upper phase of the obtained extract was used to measure the activity of the polyphenol oxidase enzyme (Lio and Hang, 2000). After measuring and calculating the mentioned traits, the obtained results were analyzed. Analysis of variance and comparison of means was done with the help of SAS software. Excel software was used to draw the figures. The mean comparison of traits was done using Duncan's test at the probability level of 5%.

3. Result and Discussion

The results of analysis of variance showed that the effect of year on flavonoid yield was significant at 1% level (P≤0.01), but it had no significant effect on phenol. The effect of fertilizer on phenol and flavonoids was significant at 1% level (P≤0.01). The effect of density on phenolic and flavonoid traits was also significant at 1% level (P≤0.01). The interaction effect of year and fertilizer on phenol and flavonoid was significant at 1% level (P≤0.01). The interaction effect of year and density, the interaction effect of fertilizer and density, and the interaction effect of year and fertilizer and density were not significant. The results of analysis of variance showed that the effect of year on the enzyme ascorbate peroxidase, catalase, and polyphenol oxidase and superoxide dismutase was significant at 1% level

($P \leq 0.01$). The effect of fertilizer on all enzymes of ascorbate peroxidase, catalase, and polyphenol oxidase and superoxide dismutase was significant at 1% level ($P \leq 0.01$). The effect of condensation on ascorbate peroxidase, superoxide dismutase, polyphenol oxidase and catalase were also significant at 1% level ($P \leq 0.01$). The interaction effect of year and fertilizer on ascorbate peroxidase and superoxide dismutase was significant at 1% level ($P \leq 0.01$) and not significant on catalase and polyphenol oxidase. The interaction effect of year and density on catalase and ascorbate peroxidase was significant at 1% level ($P \leq 0.01$) and it had no significant effect on polyphenol oxidase and superoxide dismutase. The interaction effect of density and fertilizer was significant only on ascorbate peroxidase at 1% level ($P \leq 0.01$) and had no significant effect on catalase, polyphenol oxidase and superoxide dismutase. The interaction effect of year, fertilizer and density on the enzyme ascorbate peroxidase, catalase, and polyphenol oxidase and superoxide dismutase had no significant effect (Table-3).

As shown in Figure-1. The highest amount of phenol was observed in the treatment containing cow manure. Akbarpour et al. (2015) stated that it seems that the antioxidant capacity of the plant increases with the increase in the amount of total phenol and flavonoids during the use of organic fertilizers, he also stated the maximum yield of anthocyanin and flavonoid of leaves and roots and phenol of the whole flower was related to the treatment of 30 tons of manure per hectare. Based on the results of mean comparison, with increasing planting density, the amount of plant phenol increased significantly. Phenolic compounds have antioxidant properties that prevent the occurrence of oxidative stress or reduce its effects in plant cells. Flavonoids and anthocyanins are among the secondary metabolites resulting from the phenylpropanoid pathway, and the enzyme phenylalanine ammonialase is the initiator of this pathway, which converts L-phenylalanine into trans-cinnamic acid by deamination. This path is the path of biosynthesis of secondary metabolites in the cell. This key enzyme plays an essential role in the formation of phenolic compounds and is considered as one of the indicators sensitive to environmental changes (such as changes in planting density) and also one of the biochemical indicators of plant defense against environmental stress (Voget, 2010). Therefore, it seems that plant cultivation in optimal plant density is effective in enzyme synthesis, and in this way, the production of phenolic compounds, flavonoids is affected, and finally, the quantitative and qualitative performance of the plant is improved. The results of analysis of variance (Table-3) showed that the interaction effect of year and fertilizer on the phenol of the savory plant was significant. The results of mean comparison showed (Fig.1) that in the

second crop year, cow manure had a better performance than other treatments.




Fig1. The results of the comparison of the mean interaction effect of year and fertilizer on the phenolic compounds of Sahandi Savory plant

The results of analysis of variance (Table-3) showed that the effect of year as well as different fertilizers and planting density on the flavonoid compounds of Sahandi Savory plant was significant. The results of mean comparison showed that in the second cropping year, the amount of plant flavonoid compounds was higher than the first cropping year (Figure-2). The results of comparing the average effect of different fertilizers are shown in Fig.2.




Fig2. The results of comparing the mean interaction effect of year and fertilizer on the flavonoid compounds of Sahandi Savory plant

The highest amount of flavonoid was observed in the treatment containing cow manure. The results of mean comparison showed (Fig.2) that in high and medium planting density, compared to low density, the amount of flavonoid compounds of the plant increased significantly. These compounds accumulate in the vacuoles of leaf epidermal cells and work closely with a vacuolar peroxidase enzyme to detoxify oxygenated water from other cell parts or cells (Yamasaki et al., 1997). Asadi Sanam et al. (2015) stated in a research that the low density of seven plants per square meter led to an increase in root dry matter production and a decrease in the amount of chicory acid, total phenol and flavonoid content, and the antioxidant capacity of the roots of red roses. The results of analysis of variance (Table-3) showed that the interaction effect of year and fertilizer was also significant on the flavonoid compounds of the

Table 3. The results of variance analysis of phenol, flavonoid and antioxidant enzymes of sahandi Savory plant.

Sources of changes	DF	TPC	TFC	APX	SOD	CAT	PPO
Year	1	ns 6.897	** 55.206	** 1988.972	** 32.666	** 7.141	** 0.007
Error (year)	2	NS 9.955	ns 0.968	Ns 1978.97	ns 0.166	ns 0.003	ns 0.00002
Fertilizer	2	** 339.676	** 10.583	** 9551.779	** 233.185	** 1.726	** 0.006
Year x Fertilizer	2	** 20.678	** 3.350	** 14577.962	** 20.222	ns 0.090	ns 0.0002
error (year x fertilizer)	8	ns 2.657	ns 0.267	Ns 17.41	ns 1.203	ns 0.013	ns 0.00019
Density	2	** 26.423	** 2.100	** 3468.991	** 41.129	** 0.987	** 0.0015
Year x density	2	ns 5.664	ns 0.483	** 5354.678	ns 4.166	** 0.335	ns 0.0004
Fertilizer x density	4	ns 1.465	ns 0.244	** 6309.165	ns 0.824	ns 0.147	ns 0.0008
Year x Fertilizer x Density	4	NS 2.400	ns 0.104	ns 2948.533	ns 1.638	ns 0.061	ns 0.00013
Error	24	1.736	0.178	13.601	2.453	0.053	0.00018
Coefficient of Variation (%)	-	4.567	5.385	4.785	4.789	4.198	8.725

Table 4. Essential oil composition of sahandi savory under manure fertilizer, Straw fertilizer and plant density (HPD: high plant density, MPD: medium plant density, LPD: low plant density) in 2017.

Compound	RI	Control			Manure application			Straw fertilizer application		
		HPD	MPD	LPD	HPD	MPD	LPD	HPD	MPD	LPD
α -Thujene	934.61	1.11	1.05	0.54	0.88	0.68	0.54	0.51	0.68	0.53
α -Pinene	946.09	0.75	0.78	0.57	0.65	0.57	0.53	0.31	0.51	0.52
Myrcene	976.79	1.84	1.65	1.15	1.43	1.26	1.25	1.30	1.40	1.16
α -Terpinene	1043.62	1.38	0.88	0.50	1.34	1.16	0.68	0.77	0.73	0.56
p-Cymene	1053.41	47.57	56.37	48.68	43.13	44.36	48.55	26.92	50.81	50.26
γ -Terpinene	1084.13	13.22	8.46	6.21	14.75	12.71	7.41	12.25	0.13	6.34
Cis-Sabinene hydrate	1100.00	0.18	0.25	0.35	0.19	0.24	0.23	0.59	0.43	0.46
Thymol	1320.15	26.53	23.48	31.70	27.13	30.95	32.26	36.39	30.38	28.89
Carvacrol	1330.04	1.12	1.02	1.52	3.27	1.11	1.80	1.35	1.39	1.77
Thymol acetate	1363.79	0.39	0.16	0.46	0.40	0.44	0.49	0.68	0.28	0.96
E - Caryophyllene	1476.54	0.85	0.78	0.70	0.92	0.78	0.39	1.41	0.56	0.59
Spathulenol	1649.09	0.15	0.17	0.36	0.31	0.44	0.37	0.33	0.31	0.87
Caryophyllene Oxide	1659.41	0.55	0.86	1.55	1.08	1.17	1.07	1.26	1.31	2.41

Table 5. Essential oil composition of sahandi savory under manure fertilizer, Straw fertilizer and plant density (HPD: high plant density, MPD: medium plant density, LPD: low plant density) in 2018.

Compound	RI	Control			Manure application			Straw fertilizer application		
		HPD	MPD	LPD	HPD	MPD	LPD	HPD	MPD	LPD
α -Thujene	937.56	1.36	1.08	1.00	1.08	0.99	0.90	0.93	0.88	0.68
α -Pinene	946.29	0.98	0.71	0.76	0.76	0.76	0.69	0.66	0.64	0.56
Myrcene	1010.34	0.30	0.21	0.26	0.26	0.27	0.25	0.21	0.22	0.18
α -Terpinene	1045.05	1.79	1.68	1.45	1.45	1.97	1.22	1.35	1.36	1.76
p-Cymene	1054.08	47.49	43.49	45.74	48.74	36.03	42.09	44.50	42.38	29.87
γ -Terpinene	1084.53	18.84	18.23	15.32	15.32	22.01	14.03	14.74	15.59	20.74
Cis-Sabinene hydrate	1100.00	0.14	0.16	0.16	0.16	0.15	0.22	0.24	0.29	0.31
Thymol	1322.00	21.59	26.05	24.04	24.04	29.86	31.14	28.75	30.37	36.56
Carvacrol	1326.82	1.11	1.06	0.97	0.97	0.82	1.29	1.16	1.18	1.05
Thymol acetate	364.32	0.88	0.98	0.62	0.62	1.07	0.88	0.86	1.60	2.77
E - Caryophyllene	1480.12	0.69	0.75	0.77	0.77	0.85	0.84	0.76	0.95	1.10
Spathulenol	1654.17	0.51	0.28	0.34	0.34	0.37	0.31	0.33	0.50	0.54
Caryophyllene Oxide	166.11	0.47	0.69	0.80	0.80	0.62	0.89	0.90	0.97	1.04

savory plant. Based on the results of mean comparison, cow manure caused a significant difference in plant flavonoid compounds in two crop years. The highest amount of flavonoid was observed in the treatments of the second year and cow manure (Fig.2).

Acknowledgments

This work was supported by the, shar-e-Qods Branch, Islamic Azad University, Tehran, Iran and Research Institute of Forests and Rangelands, Tehran, Iran.

References

- Akbarinia, A., Sefidkon, F., & Razaz Hashemi, S.R. (2009). Essential oil components of cultivated and wild accessions of *Satureja sahendica* Bornm. *Iranian Journal of Medicinal and Aromatic Plants*, 25(3), 376-385.
- Akbarpour, A., Kavooosi, B., Hosseinifarahi, M., Tahmasebi, S., & Gholipour, S. (2015). Evaluation of Yield and Phytochemical Content of Different Iranian Garlic (*Allium sativum* L.) Ecotypes. *International Journal of Horticultural Science and Technology*, 8(4), 385-400.
- Akintoye, H.A., Lucas, E.O., & Kling, J. (1997). Effects of density of planting and time of nitrogen application on maize varieties in different ecological zones of West Africa. *Soil Science and Plant Analysis*, 28, 1163-1175.
- Batra, P., & Sharma, A.K. (2013). Anti-cancer potential of flavonoids: recent trends and future perspectives. *Biotechnology*, 3(6), 439-459.
- Borsani, O., Valpuesta, V., & Bottela, A.M. (2001). Evidence for a role of salicylic acid in the oxidative damage generated by NaCl and osmotic stress in *Arabidopsis* seedlings. *Plant Physiology*, 126, 1024-1030.
- Brunetti, C., DiFerdinando, M., Fini, A., Pollastri, S., & Tattini, M. (2013). Flavonoids as antioxidants and developmental regulators: relative significance in plants and humans. *International Journal of Molecular Science*, 14, 3540-3555.
- Chang, C., Yang, M., Wen, H., & Chern, J. (2002). Estimation of total flavonoid content in propolis by two complementary colorimetric methods. *Journal of Food and Drug Analysis*, 10, 178-182.
- Donat, M.G., Lowry, A.L., Alexander, L.V., O'Gorman, P.A., & Maher, N. (2016). More extreme precipitation in the world's dry and wet regions. *Nature and Climate Change*, 6, 508-513, <https://doi.org/10.1038/Nclimate2941>, 2016.
- Dreosti, I.E. (2000). Antioxidant polyphenols in tea, cocoa, and wine. *Nutrition*, 16(7-8), 692-694. doi: 10.1016/s0899-9007(00)00304-x. PMID: 10906600.
- Feller, C., Blanchart, E., Bernoux, M., Lai, R., & Manlay, R. (2012). Soil fertility concepts over the past two centuries: the importance attributed to soil organic matter in developed and developing countries. *Archives of agronomy and Soil Science*, 58(S1), S3-S21.
- Gopčević, K., Grujić, S., Arsenijević, J., Karadžić, I., Izrael-Živković, L., & Maksimović, Z. (2019). Phytochemical Properties of *Satureja kitaibelii*, Potential Natural Antioxidants: a New Insight, p1-7.
- Gozubenli, H., Sener, O., Konuskan, O., & Kilinc, M. (2003). Effect of hybrid and plant density on grain yield and yield components of maize (*Zea mays* L.). *Indian Journal of Agronomy*, 48, 203-205.
- Hajhashemi, V., Ghannadi, A., & Pezeshkian, S.K. (2002). Anti-nociceptive and anti-inflammatory effects of *Satureja hortensis* L. extracts and essential oil. *Journal of Ethnopharmacology*, 82, 83-87.
- Harborne, J.B., Turner, B.L. (1984). *Plant Chemosystematics*, Academic Press, London, 123.
- Lovelli, S. (2019). Dryland farming and the agronomic management of crops in arid environments. *Journal of Agronomy*, 18, 49-54.
- Marinova, D., Ribarova, F., & Atanassova, M. (2005). Total Phenolics and Total Flavonoids in Bulgarian Fruits and Vegetables. *Journal of the University of Chemical Technology and Metallurgy*, 40(3) 3, 255-260.
- Mohebi, A. (2016). The Effect of Integrated Soil Fertility and Nutrition Management on Quantitative and Qualitative Characteristics of Date Fruit, *Journal of Beh Zeraee Agriculture*, 18, 851-860.
- Omidbaigi, R. (2005). Production and Processing of Medicinal Plants. Astane Ghodse Razavi, Mashhad, Iran, p 347.
- Omidbaygi, R. (1995). Approaches for Production and Processing of Medicinal Plants (Vol. I). Tarrahan Nashr Publication, Tehran, Iran, 424p.
- Ouchikh, O., Chahed, T., Ksouri, R., Ben Taarit, M., Faleh, H., Abdelly, C., Kchouk, M.E., & Marzouk, B. (2011). The Effects of Extraction Method on the Measured Tocopherol Level and Antioxidant Activity of *L. nobilis* Vegetative Organs. *Journal of Food Composition and Analysis*, 24, 103-110. <https://doi.org/10.1016/j.jfca.2010.04.006>
- Pistelli, L., & Giorgi, I. (2012). Antimicrobial properties of flavonoids. In: Patra AK (ed) *Dietary phytochemicals and microbes*. Springer Science+Business Media, Dordrecht, pp 33-91.

- Stewart, B.A. (2016a). Dryland Farming. In: Reference Module in Food Science, Smithers, G.W. (Ed.), Elsevier, New York, USA.
- Stewart, B.A., & Thapa, S. (2016b). Dryland Farming: Concept, Origin and Brief History. In: Innovations in Dryland Agriculture, Farooq, M. and K.H.M. Siddique (Eds.), Springer, Cham, Switzerland, pp: 3-29.
- Voget, T. (2010). Phenylpropanoid biosynthesis. *Molecular Plant*, 3, 2-20
- Yamasaki, H., Sakihama, Y., & Ikehara, N. (1997). Flavonoid–Peroxidase reaction as a detoxification mechanism of plant cells against H₂O₂. *Plant Physiology*, 115, 1405-1412.
- Yang, J., Reichert, P., Abbaspour, K.C., & Yang, H. (2007). Hydrological modelling of the Chaohe Basin in China: Statistical model formulation and Bayesian inference. *Journal of Hydrology*, 340, 167–182.