

The Effect of Propolis Extracts on Healthy Vascular Aging Mechanism by The Telomere Regulation in Huvec Cells

Seyda Kaya¹, Sevgi Durna Daştan^{2,3}, Zeliha Selamoglu^{4,5,6*}, Taner Dastan⁷, Ainash Oshibayeva⁸, Yerbol Baltabay⁴

¹Department of Medical Services and Techniques, Medical Laboratory Techniques Program, Vocational School of Health Services, Gaziantep Islam Science and Technology University, Gaziantep / Türkiye

ORCID: <https://orcid.org/0000-0001-8489-8687>

E-mail: sydkaya58@gmail.com

²Department of Biology, Faculty of Science, Sivas Cumhuriyet University, Sivas, Türkiye

³Department of Beekeeping Development, Application and Research Center, Sivas Cumhuriyet University, Sivas, Türkiye

ORCID: <https://orcid.org/0000-0003-4946-5602>

E-mail: sdurna@cumhuriyet.edu.tr

⁴Department of Biology, Faculty of Sciences, Khoja Akhmet Yassawi International Kazakh-Turkish University, Turkestan, Kazakhstan

⁵Department of Biology, Faculty of Sciences, Western Caspian University, Baku, Azerbaijan

⁶Department of Medical Biology, Medicine Faculty, Nigde Omer Halisdemir University, Nigde, Türkiye

ORCID: <https://orcid.org/0000-0001-9056-6435>

*Corresponding Author's E-mail: zselamoglu@ohu.edu.tr

⁷Department of Biochemistry, Faculty of Science, Sivas Cumhuriyet University, Sivas, Türkiye

ORCID: <https://orcid.org/0000-0003-0296-6979>

E-Mail: tdastan@cumhuriyet.edu.tr

⁸Department of Public Health and Scientific Research, Faculty of Medicine, Khoja Akhmet Yassawi International Kazakh-Turkish University, Turkestan, Kazakhstan

ORCID: <https://orcid.org/0000-0002-5655-5465>

E-mail: ainash.oshibayeva@ayu.edu.kz

⁴Department of Biology, Faculty of Sciences, Khoja Akhmet Yassawi International Kazakh-Turkish University, Turkestan, Kazakhstan

ORCID: <https://orcid.org/0009-0003-6963-9873>

E-mail: yerbol.baltabay@bk.ru

ABSTRACT

Introduction: Propolis is a very important natural bioactive product with potent antioxidant properties; however, the effects of Kazakhstan propolis on telomere-associated molecular pathways remain largely unexplored. The objective was to comparatively evaluate the antioxidant activity, phytochemical composition, cytotoxicity, and telomere-related gene expression profiles of different solvent extracts obtained from propolis collected from four regions of southern Kazakhstan.

Methods: Ethanolic, methanolic, water, and ethyl acetate extracts of four propolis samples (Pro1–Pro4) were prepared. Total phenolic content, 2,2-Diphenyl-1-picrylhydrazyl (DPPH) radical scavenging activity, total flavonoid content, and HUVEC cell viability were determined. Relative expression levels of *TERT*, *TERF1*, *TERF2*, and *PINX1* were quantified by qRT-PCR using the fold change $\Delta\Delta C_t$ method with *GAPDH*.

Results: Methanolic extracts of samples generally showed the highest phenolic and flavonoid contents and antioxidant activity. All extracts showed low cytotoxicity toward HUVEC cells, indicating good cellular compatibility. Gene expression analysis demonstrated that methanol extracts markedly enhanced *TERF1* and *TERT* expression, particularly in the Pro2 and Pro3 samples, whereas ethanol extracts produced intermediate responses, and aqueous extracts showed comparatively weaker effects. *TERF2* expression exhibited moderate upregulation, while *PINX1* showed variable but non-significant alterations among treatment groups. One-way ANOVA revealed significant differences for *TERF1*, *TERT*, and *TERF2*.

Conclusions: These data present that phenolic-rich propolis extracts may support to endothelial health by regulating oxidative damage with telomere homeostasis, and highlight Kazakhstan propolis as a healthy natural material of important compounds for healthy vascular aging.

Keywords: Anti-aging, Antioxidant activity, Endothelial cells, HUVEC, Kazakhstan propolis, qRT-PCR, Telomere.

1. Introduction

The global increase in life expectancy has been accompanied by a rise in age-related disorders, making healthy aging a major biomedical challenge. Among the mechanisms underlying aging, oxidative stress is a principal driver of endothelial dysfunction, chronic inflammation, genomic instability, and telomere shortening. Produced reactive oxygen species bring redox homeostasis, damage proteins, lipids, and nucleic acids. Because vascular endothelial cells are continuously exposed to circulating oxidants and inflammatory mediators, they are particularly susceptible to oxidative injury. Consequently, preservation of endothelial integrity is an important target for preventing cardiovascular and age-associated diseases [1].

Telomeres are nucleoprotein molecules at chromosome peripheral ends that preserve genomic stability during cell division. Progressive telomere shortening is a hallmark of cellular aging and is strongly influenced by oxidative stress. Oxidative lesions within telomeric regions are repaired less efficiently, accelerating telomere attrition and premature senescence. Telomere integrity depends on coordinated regulation of telomeric repeat-binding factor 1 (*TERF1*), telomeric repeat-binding factor 2 (*TERF2*), telomerase reverse transcriptase (*TERT*), and *PIN2/TERF1*-interacting telomerase inhibitor 1 (*PINX1*). These genes are associated with endothelial dysfunction, impaired tissue regeneration, and chronic diseases. Thus, strategies that reduce oxidative stress while maintaining telomere homeostasis have attracted increasing attention as potential interventions against cellular aging. The telomere shortening triggers apoptosis or senescence [2]. Telomerase is the principal enzyme responsible for maintaining telomere length [3] and is a ribonucleoprotein complex composed of the telomerase RNA component (TERC) and catalytic reverse transcriptase subunit (*TERT*) [4]. Although telomerase activity is minimal in most normal somatic cells, it is highly active in stem and cancer cells, highlighting its role in cellular aging and proliferation [5].

Natural products are rich sources of bioactive molecules targeting oxidative stress-related pathways. Propolis has gained considerable attention because of its diverse phytochemical composition and broad biological activities. Plant resins mixed with beeswax and salivary enzymes by honey bees, propolis contains more than 500 identified compounds [6], contributing to its cardioprotective, anti-inflammatory, antioxidant, antimicrobial, immunomodulatory, antiviral, and anticancer properties. Many of these effects involve oxidative stress-sensitive pathways, including *Nrf2*, *NF-κB*, *MAPK* [1,7]. Its protective effects on vascular endothelium are particularly noteworthy. Propolis has been shown to reduce

intracellular ROS and lipid peroxidation, preserve nitric oxide bioavailability, inhibit endothelial apoptosis, and attenuate inflammatory signaling, suggesting effects on both oxidative balance and cellular longevity [8,9].

Propolis activity varies considerably with geographical origin, local flora, climatic conditions, bee species, harvesting season, and extraction methodology. Different solvents influence the phenolic contents, resulting in distinct phytochemical profiles and antioxidant and cytoprotective potentials. Comparative evaluation of solvent-dependent extracts is therefore essential for identifying biologically active preparations and understanding their mechanisms. Most previous studies have focused on antioxidant or antimicrobial activity, whereas less attention has been given to endothelial biology and telomere-associated responses. Thus, the potential effects of Kazakhstan propolis on telomere regulatory pathways remain largely unknown.

HUVECs are widely used to investigate endothelial physiology, oxidative stress, vascular aging, and cytoprotective mechanisms. Because endothelial dysfunction is an early event in numerous chronic diseases, HUVECs provide an appropriate model for evaluating natural compounds. Simultaneous assessment of antioxidant activity, cytotoxicity, cellular morphology, and telomere-associated gene expression provides a more comprehensive evaluation of endothelial protection than antioxidant assays alone. Endothelial aging is a major risk factor for cardiovascular disease, while telomere shortening is a key biomarker of this process [10]. Therefore, HUVECs provide a suitable in vitro system for investigating telomere-related responses.

Accordingly, this study comparatively investigated propolis samples collected from four geographically distinct regions of southern Kazakhstan. This study aims to present new data for protecting effects on endothelial structure of Kazakhstan propolis and its possible role in oxidative stress-mediated telomere homeostasis, supporting its potential as a natural intervention for healthy aging and vascular protection.

2. Materials & Methods

2.1. Propolis Samples and Extraction

Samples were obtained from four geographically distinct locations in southern Kazakhstan (Table 1). Later 5 g of each sample was added with 100 mL of methanol, ethanol, ethyl acetate,

and distilled water by separately and incubated with shaking for 48 h in RT. After that they were filtrated through Whatman No. 1 paper. The rotary evaporator at 40°C for organic solvent extracts, whereas freeze-dried were aqueous extracts to concentrate the extracts.

Table 1. Propolis samples collected from Kazakhstan

Samples	Location
Pro1	Station 5: Zhanakorgan district
Pro2	Station 5: Kyzylorda district
Pro3	Station 5: Otrar district
Pro4	Station 5: Shieli district

2.2. DPPH Radical Scavenging Assay

Antioxidant activity was assessed using the DPPH assay as previously defined [11]. Sample solutions (40 μ L) prepared in water were mixed with 160 μ L of DPPH solution (0.4 mM) dissolved in ethanol. The reaction mixture was incubated in the dark at room temperature for 30 min. Water was used as the negative control, while BHA (Butylated hydroxyanisole) was used as standard antioxidants. Absorbance was collected at 517 nm. The percentage inhibition were calculated from absorbance values, and IC₅₀ values were calculated.

2.3. Total phenolic content Assay

TPC was determined using the Folin–Ciocalteu colorimetric assay with minor modifications to the method described by Clarke et al. [11]. Briefly, 20 μ L of extract solution was mixed with 100 μ L of 10% Folin–Ciocalteu reagent and incubated for 5 min. Subsequently, 80 μ L of 7.5% sodium carbonate solution was added and the reaction mixture was incubated in the dark for 30 min at room temperature. Absorbance was measured at 765 nm using a BioTek Epoch microplate spectrophotometer. Gallic acid was used for preparation of the calibration curve and results were expressed as mg gallic acid equivalents (GAE)/g dry extract.

2.4. Total flavonoid content Assay

TFC was quantified using the aluminum chloride colorimetric method [12]. Extracts were reacted with 2% aluminum chloride solution and incubated at room temperature for 30 min.

Absorbance was recorded at 415 nm, and TFC was expressed as mg quercetin equivalents (QE)/g extract using a quercetin calibration curve.

2.5. HUVEC Cell Culture and MTT Cytotoxicity Assay

HUVECs were cultured in DMEM containing 10% FBS and 1% penicillin–streptomycin at 37°C and 5% CO₂, subcultured with Trypsin–EDTA, and seeded into 96-well plates. Cytotoxicity was assessed by the MTT assay. Propolis extracts prepared in 1% DMSO-containing culture medium were applied at 0.01–100 mg/mL for 24 or 48 h (n = 5). Cell viability was calculated with the Vybrant MTT Cell Proliferation Assay Kit (Invitrogen) by using the manufacturer's instructions. Following 4 h incubation at 37°C, absorbance was measured at 540 nm using an Epoch microplate reader (BioTek). Viability was expressed relative to untreated controls, and IC₅₀ values were calculated using GraphPad Prism.

2.6. Morphological Assessment of HUVEC Cells

HUVEC cells were treated with propolis extracts (10 mg/mL) for 24 or 48 h. Following treatment, cellular morphology was examined using an inverted microscope (Zeiss Axiovert A1, Carl Zeiss, Germany), and images were acquired with ZEN 3.1 (ZEN lite) imaging software (Carl Zeiss).

2.7. RNA Isolation

HUVEC cells were treated with propolis extract for 24 or 48 h prior to RNA isolation. Total RNA was extracted using Hydrozol reagent according to the manufacturer's protocol (A.B.T.TM RNA Purification Kit). Briefly, cell lysates were subjected to chloroform phase separation. The purified RNA pellets were dissolved in RNase-free water.

2.8 cDNA Synthesis

Complementary DNA (cDNA) was synthesized from total RNA using the GLPBIO SuperFast RT Master Mix for qPCR kit with gDNA Remover (Cat. No. GK10031) according to the manufacturer's instructions. Briefly, RNA (0.01–1 µg) was first treated with 2 µL of 5× gDNA Removal Mix in a final volume of 10 µL at 42°C for 2 min. Subsequently, 4 µL of 5× SuperFast RT Mix and 6 µL of DEPC-treated ddH₂O were added to obtain a final reaction volume of 20 µL, followed by reverse transcription at 50°C for 5 min.

2.9. Quantitative Real-Time PCR (qRT-PCR)

Gene expression values were presented using synthesized cDNA and gene-specific primers targeting *GAPDH*, *TERF1*, *TERF2*, *TERT*, and *PINX1* (Table 2). *GAPDH* was used as the house keeping gene. Each PCR reaction consisted of 2.5 μ L BlasTaq™ 2 $\times$ qPCR Master Mix, 0.5 μ L each of forward and reverse primers (1 μ M), 0.5 μ L template cDNA (50 ng), and 1.5 μ L nuclease-free water, giving a final volume of 5.5 μ L. Amplification was carried out using an Applied Biosystems Real-Time PCR System with an initial enzyme activation at 95°C for 3 min, followed by 40 cycles of denaturation at 95°C for 15 s and annealing/extension at 60°C for 1 min. Using threshold cycle (Ct) values for $\Delta\Delta$ Ct method, the fold change results were visualized graphically.

Table 2. Primers used for qRT-PCR analysis

Full gene name	Gene primers	Primer sequence (5'→3')
Glyceraldehyde-3-phosphate dehydrogenase	GAPDH-F	5'-GTCTCCTCTGACTTCAACAGCG-3'
	GAPDH-R	5'-ACCACCCTGTTGCTGTAGCCAA-3'
Telomeric Repeat Binding Factor 1	TERF 1-F	5'-CATGGAACCCAGCAACAAGACC-3'
	TERF 1-R	5'-CTGCTTTCAGTGGCTCTTCTGC-3'
Telomeric Repeat Binding Factor 2	TERF 2-F	5'-GTGGAAAAGCCACCCAGAGAAC-3'
	TERF 2-R	5'-TGCAAAGGCTGCCTCAGAATCC-3'
Telomerase Reverse Transcriptase	TERT-F	5'-GCCGATTGTGAACATGGACTACG-3'
	TERT-R	5'-GCTCGTAGTTGAGCACGCTGAA-3'
PIN2/TERF1-Interacting Telomerase Inhibitor 1	PINX 1-F	5'-TCCAGAGGAGAACGAAACCACG-3'
	PINX 1-R	5'-TACGTTCCACCTGCGTCTCAGA-3'

2.10. Statistical analyses

The statistics were performed using GraphPad Prism 10.0. Relative gene expression ($\Delta\Delta$ Ct values) was analyzed by one-way analysis of variance (ANOVA), followed by Tukey's multiple comparison test. Data are presented as mean $\pm$ standard deviation (SD). Differences were considered statistically significant at $p < 0.05$.

3. Results

3.1. Extraction Yield of Propolis

Extract yields were calculated as percentages (Table 3) and the yield values were between 6-12%. Total phenolic content was determined because phenolic compounds are closely associated with the biological activity of propolis extracts and may vary according to geographical and seasonal conditions.

3.2. DPPH Radical Scavenging Activity

The antioxidant activity of propolis extracts are presented in Table 3. IC₅₀ values ranged from 97.5 to 213.2 µg/mL, with lower values indicating stronger activity. Overall, the extracts showed moderate-to-high antioxidant activity compared with BHA.

3.3. Total Phenolic and Flavonoid Content

Total phenolic content ranged from 99.4 to 216.3 mg GAE/g extract, while total flavonoid content ranged from 6.9 to 13.6 mg QE/g extract (Table 3).

3.4. Cell Culture Findings

Propolis extracts were prepared at 100 mg/mL and diluted to 0.01–100 mg/mL. Based on MTT and microscopic findings, 10 mg/mL was selected for subsequent experiments. Cell morphology was evaluated after 24 and 48 h using a Zeiss Axio inverted microscope at 10× magnification (Figure 1). Propolis-treated HUVECs maintained normal morphology and proliferation without evident cellular stress. Given the IC₅₀ values of 5.6–25.4 mg/mL, the extracts showed weak cytotoxicity, and no further anticancer analyses were performed. Experiments were conducted in five independent replicates (n = 5).

230

231 **Table 3.** Percentage yields, DPPH activity, TPC and TFC content of propolis extracts

Samples	Extracts	Yield %	DPPH• radical scavenging activity IC50 (µg/mL)	Total Phenolic Content (mg GAE/g extract ±SD)	Total Flavonoid Content (mg QE/g extract ±SD)	Cell Culture MTT results (IC ₅₀ (mg/mL))
Pro1	Water	7%	195.3 ± 6.9	146.2 ± 5.6	7.9 ± 0.6	18.7
	Methanol	10%	148.79 ± 3.6	178.5 ± 8.6	9.9 ± 1.3	10.5
	Ethanol	11%	136.9 ± 4.1	173.4 ± 8.9	10.3 ± 1.6	7.5
	Ethyl acetate	8%	165.89 ± 6.3	161.7 ± 7.9	8.3 ± 1.0	19.5
Pro2	Water	6%	189.63 ± 6.9	153.1 ± 6.6	6.9 ± 0.5	20.2
	Methanol	9%	138.5 ± 6.9	179.8 ± 7.6	10.5 ± 0.9	10.1
	Ethanol	10%	135.9 ± 3.6	173.2 ± 7.3	11.9 ± 1.3	6.3
	Ethyl acetate	8%	154.6 ± 9.3	160.6 ± 6.9	8.3 ± 1.0	17.6
Pro3	Water	7%	169.5 ± 3.6	158.3 ± 6.9	7.1 ± 1.1	19.6
	Methanol	11%	141.3 ± 9.7	181.1 ± 7.6	12.4 ± 1.3	9.6
	Ethanol	11%	148.9 ± 6.3	196.7 ± 9.9	13.6 ± 1.1	5.6
	Ethyl acetate	9%	158.6 ± 9.3	161.7 ± 5.6	8.7 ± 1.0	16.3
Pro4	Water	7%	213.2 ± 6.9	99.4 ± 3.3	8.4 ± 0.9	25.4
	Methanol	10%	144.9 ± 6.3	126.7 ± 5.9	12.5 ± 0.8	9.1
	Ethanol	12%	97.5 ± 6.3	216.3 ± 8.6	12.3 ± 1.3	7.9
	Ethyl acetate	10%	118.6 ± 3.9	143.3 ± 5.6	7.7 ± 2.6	15.9
BHA: Reference substance for DPPH		-	10.1 ± 1.95	-	-	

232




Figure 1. Representative microscopic images of control and propolis-treated HUVEC cells after 24 h of incubation. Images were obtained using a Zeiss Axio inverted microscope

3.5. Effects of Different Propolis Extracts on Telomere-Related Gene Expression in HUVEC Cells

Following RNA quality assessment and cDNA synthesis, the relative expression of *TERT*, *TERF1*, *TERF2*, and *PINX1* was quantified by qRT-PCR using *GAPDH* as the reference gene and untreated cells as the calibrator. Propolis treatment altered telomere-related gene expression according to sample origin and extraction solvent (Table 4). *TERF1* showed the strongest response, particularly in methanolic extracts, with Pro3-Methanol producing the highest expression, followed by Pro2- and Pro4-Methanol. Methanolic extracts induced approximately 6–16-fold higher *TERF1* expression than the control (Table 4, Figure 2). *TERT* expression was highest in Pro3-Methanol and Pro4-Methanol, while *PINX1* showed moderate and *TERF2* comparatively modest induction, with Pro3-Methanol producing the strongest responses.

Extraction solvent had a greater effect than geographical origin, with the overall pattern Methanol > Ethanol > Ethyl acetate > Water; Pro3 showed the strongest and Pro1 the weakest response. One-way ANOVA revealed significant differences for *TERF1* ($F = 8.404$, $p < 0.0001$), *TERT* ($F = 5.670$, $p < 0.0001$), and *TERF2* ($F = 4.363$, $p < 0.001$), but not *PINX1* ($F = 1.214$, $p = 0.295$) (Table 5, Figure 2). Tukey's test showed significantly higher *TERT* expression in several Pro3, Pro2, Pro1, and Pro4 groups ($p < 0.05$), while *TERF1* upregulation was significant only

for Pro2- and Pro3-Methanol extracts. No significant pairwise differences were detected for *TERF2* or *PINX1* after correction (Table 4, Figure 2).

Table 4. Relative expression levels of telomere-associated genes in HUVEC cells treated with different propolis extracts (The statistical difference is significant at * $p < 0.05$; ** $p < 0.01$ and *** $p < 0.001$)

<i>TERT</i> gene expression values and Tukey HSD compare				
Sample	Water	Ethanol	Methanol	Ethyl acetate
Propolis 1	1.48 ± 0.22	$1.62 \pm 0.19^{**}$	2.61 ± 0.28	1.86 ± 0.20
Propolis 2	$1.34 \pm 0.17^{**}$	2.08 ± 0.23	2.29 ± 0.21	$1.77 \pm 0.18^{**}$
Propolis 3	$1.92 \pm 0.21^{***}$	$2.47 \pm 0.25^{***}$	$3.21 \pm 0.34^{***}$	2.53 ± 0.29
Propolis 4	1.86 ± 0.18	$2.35 \pm 0.24^{**}$	2.89 ± 0.31	2.41 ± 0.25
<i>TERF2</i> gene expression values and Tukey HSD compare				
Propolis 1	1.34 ± 0.18	2.24 ± 0.24	2.91 ± 0.31	1.82 ± 0.19
Propolis 2	1.05 ± 0.12	2.03 ± 0.22	1.94 ± 0.21	1.37 ± 0.15
Propolis 3	1.61 ± 0.18	2.27 ± 0.23	3.08 ± 0.33	2.06 ± 0.22
Propolis 4	0.82 ± 0.10	1.02 ± 0.13	1.26 ± 0.16	1.11 ± 0.12
<i>PINX1</i> gene expression values and Tukey HSD compare				
Propolis 1	0.81 ± 0.09	1.34 ± 0.16	1.62 ± 0.19	1.42 ± 0.17
Propolis 2	0.94 ± 0.11	1.51 ± 0.18	1.46 ± 0.17	1.13 ± 0.15
Propolis 3	1.68 ± 0.19	3.76 ± 0.35	4.62 ± 0.41	2.51 ± 0.24
Propolis 4	1.72 ± 0.20	3.04 ± 0.29	4.08 ± 0.37	2.44 ± 0.23
<i>TERF1</i> gene expression values and Tukey HSD compare				
Propolis 1	2.41 ± 0.26	4.55 ± 0.41	6.91 ± 0.54	2.78 ± 0.30
Propolis 2	4.23 ± 0.39	9.63 ± 0.81	$13.81 \pm 1.05^{***}$	5.12 ± 0.48
Propolis 3	3.76 ± 0.33	11.26 ± 0.95	15.83 ± 1.18	6.08 ± 0.52
Propolis 4	3.64 ± 0.30	7.11 ± 0.62	12.35 ± 0.94	5.41 ± 0.46

Table 5. One-way ANOVA results of genes

Gene	F value	p value	Significance
<i>TERT</i>	5.670	0.000002	Significant
<i>TERF1</i>	8.404	1.01×10^{-8}	Very Significant
<i>TERF2</i>	4.363	0.000043	Significant
<i>PINX1</i>	1.214	0.295	Not Significant




Figure 2. Relative expression levels of telomere-associated genes (*TERT*, *TERF1*, *TERF2* and *PINX1*) in HUVEC cells following 24-hour treatment with four propolis samples extracted using different solvents

4. Discussion

It is demonstrated that extraction solvent markedly changed the biological activity of extracts. Methanolic extracts showed the highest phenolic and flavonoid contents, antioxidant activity, and modulation of telomere-associated genes in HUVECs, whereas aqueous extracts generally showed weaker responses. This solvent dependence likely reflects differences in the extraction efficiency of phenolic acids. Methanol and ethanol efficiently recover compounds such as

274 chrysin, galangin, CAPE, quercetin derivatives, while water preferentially extracts highly polar
275 constituents. The association between total phenolic content and DPPH activity further supports
276 the contribution of phenolics to propolis antioxidant capacity.

277 Oxidative stress promotes telomere shortening and endothelial senescence, as guanine-rich
278 telomeric DNA is particularly susceptible to oxidative damage. Thus, the higher antioxidant
279 activity of methanolic and ethanolic extracts may partly explain their stronger effects on
280 telomere-related genes. By reducing oxidative damage, propolis antioxidants may help preserve
281 telomere integrity and regulate telomere-maintenance pathways.

282 *TERF1* showed the most pronounced response, with methanolic extracts producing the highest
283 induction across propolis origins. Pro2- and Pro3-Methanol increased *TERF1* expression by
284 approximately 6–16-fold relative to the control, and Tukey's test confirmed significant
285 upregulation in these groups. As a core shelterin component, *TERF1* binds double-stranded
286 telomeric DNA and contributes to telomere length homeostasis and chromosome-end
287 protection. Although excessive *TERF1* can restrict telomerase accessibility, moderate induction
288 may support telomere stabilization under oxidative stress.

289 The strong *TERF1* response to methanolic extracts may reflect their greater recovery of
290 flavonoids and phenolic acids, including CAPE, chrysin, galangin, pinocembrin, quercetin, and
291 caffeic acid derivatives. These compounds may enhance Nrf2-dependent antioxidant advocacy
292 and smilate oxidative stress-related NF- κ B signaling, providing a potential mechanistic link
293 between antioxidant activity and *TERF1* regulation.

294 Geographical origin had a smaller effect than extraction solvent. Although Pro3 generally
295 showed the strongest responses, all samples followed the same activity pattern (Methanol>
296 Ethanol>Ethyl acetate >Water), indicating that extraction efficiency was a major determinant
297 of biological activity.

298 *TERT* expression was also strongly affected by propolis treatment, with methanolic extracts
299 consistently producing the highest levels. Pro3-Methanol showed the strongest increase (3.21
300 ± 0.34 -fold), followed by Pro4-Methanol (2.89 ± 0.31 -fold), whereas water extracts showed
301 weaker and ethanol and ethyl acetate intermediate responses. Significant differences among
302 groups and higher expression in several Pro3-treated groups were confirmed statistically.
303 Telomerase contributes to chromosome stability by compensating for progressive telomere

shortening [3,13]. Although its expression is limited in most differentiated cells, endothelial cells retain measurable activity that contributes to vascular homeostasis. Oxidative stress can suppress *TERT* transcription, whereas antioxidant phytochemicals may restore telomerase-related signaling through *Nrf2*, *PI3K/Akt*, and *SIRT1*. Thus, increased *TERT* expression may reflect improved redox balance rather than uncontrolled proliferation.

The association between antioxidant activity and *TERT* expression further supports this interpretation, as methanolic extracts showed the lowest DPPH IC₅₀ values and highest phenolic contents. *TERF2* showed a more moderate but generally consistent increase, particularly with Pro1- and Pro3-Methanol. Although the overall group effect was significant, Tukey's analysis detected no significant pairwise differences. As a key shelterin component, *TERF2* stabilizes telomeric structures and prevents inappropriate DNA damage responses; its modest modulation may therefore reflect preservation of telomere homeostasis rather than excessive alteration of shelterin function.

In contrast, *PINX1* showed a weaker response, and overall differences among groups were not statistically significant. *PINX1* is an endogenous regulator of telomerase through its interaction with *TERT*, but it also contributes to chromosome stability and telomere architecture. Therefore, the simultaneous increase in *TERT* and *PINX1* observed in some methanol-treated groups may represent a compensatory mechanism that permits telomerase activation while preventing excessive telomere elongation. Such coordinated regulation may reflect maintenance of telomere homeostasis rather than contradictory biological effects.

Overall, extraction solvent exerted a greater influence on telomere-related gene expression than geographical origin, with the consistent hierarchy of Methanol>Ethanol>Ethyl acetate >Water. Nevertheless, considerable variation among Pro1–Pro4 was observed, which is consistent with the known influence of botanical origin, geography, climate, and local vegetation on propolis composition [14–16]. Pro3 generally produced the strongest responses, particularly for *TERF1* and *TERT*. Its activity may be related to a richer profile of flavonoids and phenolic compounds capable of modulating *PI3K/Akt*, *MAPK*, *Nrf2*, *FOXO*, and *NF-κB* pathways, which interact with telomerase regulation and cellular senescence [17].

The coordinated changes in *TERF1*, *TERT*, *TERF2*, and *PINX1* suggest that propolis does not act on a single telomere-associated target but modulates multiple components involved in telomere protection and endothelial homeostasis. Alterations in shelterin proteins may

accompany telomeric remodeling under oxidative stress, as the shelterin complex functions as an integrated regulatory network [18]. The simultaneous induction of *TERT* and *PINX1* may further reflect negative-feedback regulation following telomerase activation, supporting a balanced rather than excessive telomere response. In this case, the data from this research sign that propolis may influences as a cellular modulator, fine-tuning telomere-associated pathways according to its phytochemical composition.

Several limitations should be acknowledged. First, only mRNA expression was measured; protein abundance and enzymatic activity were not assessed. Second, telomerase activity was not directly determined using the TRAP assay; therefore, increased *TERT* expression cannot be considered definitive evidence of increased telomerase activity. Third, telomere length was not measured by qPCR or terminal restriction fragment analysis. Accordingly, the functional consequences of the observed transcriptional changes remain to be experimentally validated.

5. Conclusion

Finally propolis extracts may effect the expression of key telomere-associated genes in HUVEC cells, with methanolic extracts generally producing the strongest responses. The findings further indicate that extraction solvent had a greater influence on telomere-related gene expression than geographical origin, highlighting the importance of extraction strategy in determining the biological activity of propolis. Future studies should integrate transcriptomic, proteomic, and functional approaches, including TRAP assays, shelterin protein analyses, ROS measurements, and telomere-length determination, to clarify the mechanisms underlying propolis-mediated telomere regulation. LC-MS/MS or UHPLC-QTOF-MS profiling could further link specific bioactive compounds to telomere-related responses. Overall, these findings identify solvent-specific propolis extracts as potential modulators of cellular aging, although further mechanistic and functional validation is required before therapeutic or preventive applications can be considered.

Acknowledgment

We thank the Scientific Committee of the Ministry of Science and Higher Education of the Republic of Kazakhstan for supporting this project (**BR24992814**).

Authors' Contribution

Conceptualization: ZS, SDD. Data curation: ZS, SDD, SK, YB. Formal analysis: ZS, SK, AO. Methodology: ZS, SDD, SK, AO. Software: SK, YB, TD. Visualization: SDD, SK, TD. Investigation: ZS, SDD, SK. Writing- original draft: SDD, SK, Writing- review and editing: ZS, SDD, SK.

Ethics

Ethics committee approval is not required for this study.

Conflict of Interest

No potential conflicts of interest relevant to this article were reported.

Funding Sources

This study is supported by the project (**BR24992814**) on the development of innovative technologies and creation of scientific infrastructure for the sustainable development of the South Kazakhstan region, funded by the Scientific Committee of the Ministry of Science and Higher Education of the Republic of Kazakhstan.

Data Availability

The data that support the findings of this study are available on request from the corresponding author.

Declaration on the Use of AI

During the preparation of this manuscript, the authors used generative AI tools solely to improve linguistic clarity, grammar, with the aim of improving the figures visually and readability of the English text. All scientific content, data analysis, interpretation, and conclusions were generated, verified, and approved by the authors.

References

- [1] Silva H, Francisco R, Saraiva A, Francisco S, Carrascosa C, Raposo A. The Cardiovascular Therapeutic Potential of Propolis-A Comprehensive Review. *Biology (Basel)*. 2021; 10(1):27. [DOI:10.3390/biology10010027] [PMID:33406745]
- [2] Harley CB, Futcher AB, Greider CW. Telomeres shorten during ageing of human fibroblasts. *Nature*. 1990; 345(6274):458-60. [DOI:10.1038/345458a0] [PMID:2342578]

- 393 [3] Greider CW, Blackburn EH. Identification of a specific telomere terminal transferase
394 activity in Tetrahymena extracts. *Cell*. 1985; 43(2 Pt 1):405-13. [DOI:10.1016/0092-
395 8674(85)90170-9] [PMID:3907856]
- 396 [4] Roake CM, Artandi SE. Regulation of human telomerase in homeostasis and disease. *Nat*
397 *Rev Mol Cell Biol*. 2020; 21(7):384-97. [DOI:10.1038/s41580-020-0234-z]
- 398 [5] Kim C, Kang D, Lee EK, Lee JS. Long noncoding RNAs and RNA-binding proteins in
399 oxidative stress, cellular senescence, and age-related diseases. *Oxid Med Cell Longev*. 2017;
400 2017:2062384. [DOI:10.1155/2017/2062384] [PMID:28351025]
- 401 [6] Silici S. Bilimsel Gerçeklerle Apiterapi. Ankara: Akademisyen Kitabevi; 2020.
402 [DOI:10.37609/akya.494]
- 403 [7] Hossain R, Quispe C, Khan RA, Saikat ASM, Ray P, Ongalbek D, et al. Propolis: An update
404 on its chemistry and pharmacological applications. *Chin Med*. 2022; 17(1):100.
405 [DOI:10.1186/s13020-022-00651-2] [PMID:36028892]
- 406 [8] Zulhendri F, Chandrasekaran K, Kowacz M, Ravalia M, Kripal K, Fearnley J, et al.
407 Antiviral, Antibacterial, Antifungal, and Antiparasitic Properties of Propolis: A Review. *Foods*.
408 2021; 10(6):1360. [DOI:10.3390/foods10061360] [PMID:34204859]
- 409 [9] Malekahmadi M, Pahlavani N, Heshmati J, Clayton ZS, Beigmohammadi MT, Navashenag
410 JG, et al. Effect of propolis supplementation on oxidative stress markers: A systematic review
411 of randomized controlled trials. *J Herbal Med*. 2023; 40:100679.
412 [DOI:10.1016/j.hermed.2023.100679]
- 413 [10] Minamino T, Miyauchi H, Yoshida T, Ishida Y, Yoshida H, Komuro I. Endothelial cell
414 senescence in human atherosclerosis: role of telomere in endothelial dysfunction. *Circulation*.
415 2002; 105(13):1541-4. [DOI: 10.1161/01.cir.0000013836.85741.17] [PMID:11927519]
- 416 [11] Clarke G, Ting KN, Wiart C, Fry J. High correlation of 2,2-diphenyl-1-picrylhydrazyl
417 (DPPH) radical scavenging, ferric reducing activity potential and Total Phenolics content
418 indicates redundancy in use of all three assays to screen for antioxidant activity of extracts of
419 plants from the Malaysian rainforest. *Antioxidants (Basel)*. 2013; 2(1):1-10.
420 [DOI:10.3390/antiox2010001] [PMID:26784333]
- 421 [12] Bag GC, Grihanjali Devi P, Bhaigyab T. Assessment of total flavonoid content and
422 antioxidant activity of methanolic rhizome extract of three *Hedychium* species of Manipur
423 valley. *Int J Pharm Sci Rev Res*. 2015; 30(1):154-9.
424 <http://globalresearchonline.net/journalcontents/v30-1/28.pdf>
- 425 [13] Wang F, Stewart AJ, Kasbek C, Zhao Y, Wright WE, Price CM. Human CST has
426 independent functions during telomere duplex replication and C-strand fill-in. *Cell Rep*. 2012;
427 2(5):1096-103. [DOI:10.1016/j.celrep.2012.10.007] [PMID:23142662]
- 428 [14] Bankova V, Popova M, Trusheva B. Propolis volatile compounds: chemical diversity and
429 biological activity: a review. *Chem Cent J*. 2014; 8(1):28. [DOI:10.1186/1752-153X-8-28]
430 [PMID:24812573]

431 [15] Blackburn EH, Epel ES, Lin J. Human telomere biology: A contributory and interactive
 432 factor in aging, disease risks, and protection. *Science*. 2015; 350(6265):1193-8.
 433 [DOI:10.1126/science.aab3389] [PMID:26785477]

434 [16] Huang S, Zhang CP, Wang K, Li GQ, Hu FL. Recent advances in the chemical composition
 435 of propolis. *Molecules*. 2014; 19(12):19610-32. [DOI:10.3390/molecules191219610]
 436 [PMID:25432012]

437 [17] Reinisalo M, Kårlund A, Koskela A, Kaarniranta K, Karjalainen RO. Polyphenol
 438 Stilbenes: Molecular Mechanisms of Defense Against Oxidative Stress and Age-Related
 439 Diseases. *Oxid Med Cell Longev*. 2015; 2015:340520. [DOI:10.1155/2015/340520]
 440 [PMID:26180581]

441 [18] De Lange T. Shelterin-Mediated Telomere Protection. *Annu Rev Genet*. 2018; 52:223-47.
 442 [DOI:10.1146/annurev-genet-032918-021921] [PMID:30208292]

443

444